



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PUS7L siRNA (m): sc-152595



BACKGROUND

PUS7L (pseudouridylate synthase 7 homolog (*S. cerevisiae*)-like) is a 701 amino acid protein that belongs to the pseudouridine synthase truD family and contains one TRUD domain. The PUS7L gene is conserved in chimpanzee, canine, bovine, mouse, chicken and zebrafish, and maps to human chromosome 12q12. Encoding over 1,100 genes within 132 million bases, chromosome 12 makes up about 4.5% of the human genome. A number of skeletal deformities are linked to chromosome 12 including hypochondrogenesis, achondrogenesis and Kniest dysplasia. Noonan syndrome, which includes heart and facial developmental defects among the primary symptoms, is caused by a mutant form of PTPN11 gene product, SH-PTP2. Chromosome 12 is also home to a homeobox gene cluster which encodes crucial transcription factors for morphogenesis, and the natural killer complex gene cluster encoding C-type lectin proteins which mediate the NK cell response to MHC I interaction.

REFERENCES

1. Jamieson, C.R., van der Burgt, I., Brady, A.F., van Reen, M., Elsawi, M.M., Hol, F., Jeffery, S., Patton, M.A. and Mariman, E. 1994. Mapping a gene for Noonan syndrome to the long arm of chromosome 12. *Nat. Genet.* 8: 357-360.
2. Yang, W. and Cole, W.G. 1998. Low basal transcripts of the COL2A1 collagen gene from lymphoblasts show alternative splicing of exon 12 in the Kniest form of spondyloepiphyseal dysplasia. *Hum. Mutat.* 1: S1-S2.
3. Trowsdale, J., Barten, R., Haude, A., Stewart, C.A., Beck, S. and Wilson, M.J. 2001. The genomic context of natural killer receptor extended gene families. *Immunol. Rev.* 181: 20-38.
4. Kelley, J., Walter, L. and Trowsdale, J. 2005. Comparative genomics of natural killer cell receptor gene clusters. *PLoS Genet.* 1: 129-139.
5. van der Burgt, I. 2007. Noonan syndrome. *Orphanet J. Rare Dis.* 2: 4.
6. Krejčí, J., Uhlířová, R., Galiová, G., Kozubek, S., Smigová, J. and Bártová, E. 2009. Genome-wide reduction in H3K9 acetylation during human embryonic stem cell differentiation. *J. Cell. Physiol.* 219: 677-687.
7. SWISS-PROT/TrEMBL (Q9HOK6). World Wide Web URL:
<http://www.uniprot.org/uniprot/Q9HOK6>

CHROMOSOMAL LOCATION

Genetic locus: Pus7l (mouse) mapping to 15 E3.

PRODUCT

PUS7L siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PUS7L shRNA Plasmid (m): sc-152595-SH and PUS7L shRNA (m) Lentiviral Particles: sc-152595-V as alternate gene silencing products.

For independent verification of PUS7L (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152595A, sc-152595B and sc-152595C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PUS7L siRNA (m) is recommended for the inhibition of PUS7L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PUS7L gene expression knockdown using RT-PCR Primer: PUS7L (m)-PR: sc-152595-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.