



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



R-Spondin4 siRNA (m): sc-152620

BACKGROUND

Roof plate-specific Spondins (R-Spondins) are secreted proteins that possess a Furin-like cysteine-rich domain and are involved in regulating β -catenin function. R-Spondin4, also known as RSP04, is a 234 amino acid secreted protein that, characteristic of R-Spondins, contains one Furin-like repeat and one TSP type-1 domain. R-Spondin4 interacts with heparin and is able to activate the β -catenin signaling cascade, possibly via control of the Wnt/ β -catenin signaling pathway. Defects in the gene encoding R-Spondin4 are the cause of anonychia, an autosomal recessive condition characterized by severe hypoplasia (underdevelopment) of fingernails and toenails. R-Spondin4 exists as multiple alternatively spliced isoforms that are encoded by a gene which maps to human chromosome 20.

REFERENCES

1. Kim, K.A., Zhao, J., Andarmani, S., Kakitani, M., Oshima, T., Binnerts, M.E., Abo, A., Tomizuka, K. and Funk, W.D. 2006. R-Spondin proteins: a novel link to β -catenin activation. *Cell Cycle* 5: 23-26.
2. Blaydon, D.C., Ishii, Y., O'Toole, E.A., Unsworth, H.C., Teh, M.T., Rüschendorf, F., Sinclair, C., Hopsu-Havu, V.K., Tidman, N., Moss, C., Watson, R., de Berker, D., Wajid, M., Christiano, A.M. and Kelsell, D.P. 2006. The gene encoding R-spondin 4 (RSPO4), a secreted protein implicated in Wnt signaling, is mutated in inherited anonychia. *Nat. Genet.* 38: 1245-1247.
3. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 6105736. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Nakamura, M. and Miyachi, Y. 2008. Congenital hyponychia without RSPO4 mutation. *Acta Derm. Venereol.* 88: 511-512.
5. Chishti, M.S., Kausar, N., Rafiq, M.A., Amin, M. and Ahmad, W. 2008. A novel missense mutation in RSPO4 gene underlies autosomal recessive congenital anonychia in a consanguineous Pakistani family. *Br. J. Dermatol.* 158: 621-623.
6. Brüchle, N.O., Frank, J., Frank, V., Senderek, J., Akar, A., Koc, E., Rigopoulos, D., van Steensel, M., Zerres, K. and Bergmann, C. 2008. RSPO4 is the major gene in autosomal-recessive anonychia and mutations cluster in the furin-like cysteine-rich domains of the Wnt signaling ligand R-spondin 4. *J. Invest. Dermatol.* 128: 791-796.
7. Ishii, Y., Wajid, M., Bazzi, H., Fantauzzo, K.A., Barber, A.G., Blaydon, D.C., Nam, J.S., Yoon, J.K., Kelsell, D.P. and Christiano, A.M. 2008. Mutations in R-spondin 4 (RSPO4) underlie inherited anonychia. *J. Invest. Dermatol.* 128: 867-870.
8. Kim, K.A., Wagle, M., Tran, K., Zhan, X., Dixon, M.A., Liu, S., Gros, D., Korver, W., Yonkovich, S., Tomasevic, N., Binnerts, M. and Abo, A. 2008. R-Spondin family members regulate the Wnt pathway by a common mechanism. *Mol. Biol. Cell* 19: 2588-2596.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Rspo4 (mouse) mapping to 2 G3.

PRODUCT

R-Spondin4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see R-Spondin4 shRNA Plasmid (m): sc-152620-SH and R-Spondin4 shRNA (m) Lentiviral Particles: sc-152620-V as alternate gene silencing products.

For independent verification of R-Spondin4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152620A, sc-152620B and sc-152620C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

R-Spondin4 siRNA (m) is recommended for the inhibition of R-Spondin4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor R-Spondin4 gene expression knockdown using RT-PCR Primer: R-Spondin4 (m)-PR: sc-152620-PR (20 μ l). Annealing temperature for the primers should be 55-60°C and the extension temperature should be 68-72°C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.