



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rab 13 siRNA (m): sc-152627

BACKGROUND

The Ras-related superfamily of guanine nucleotide binding proteins, which includes the R-Ras, Rap, Rac/Rec and Rho/Rab superfamilies, exhibits 30-60% homology with Ras p21. Accumulating data suggests an important role for Rab proteins, either in endocytosis or in biosynthetic protein transport. The Rab family of small G proteins play an important role in determining the specificity of vesicular transport pathways. Rab 13 and Rab 3B localize to tight junctions in epithelial cells and to cytoplasmic vesicular structures in cells lacking tight junctions. Rab 13 can be detected in the junctional complex regions of a variety of epithelia, including intestine, kidney and liver.

REFERENCES

- Novick, P. and Brennwald, P. 1993. Friends and family: the role of the Rab GTPases in vesicular traffic. *Cell* 75: 597-601.
- Ferro-Novick, S. and Novick, P. 1993. The role of GTP-binding proteins in transport along the exocytic pathway. *Annu. Rev. Cell Biol.* 9: 575-599.
- Zahraoui, A., et al. 1994. A small Rab GTPase is distributed in cytoplasmic vesicles in non polarized cells but colocalizes with the tight junction marker ZO-1 in polarized epithelial cells. *J. Cell Biol.* 124: 101-115.
- Leek, J.P., et al. 1997. Assignment of the Rab13 gene (RAB13) to human chromosome band 12q13 by *in situ* hybridization. *Cytogenet Cell Genet.* 79: 210-211. PMID: 9605854
- Sheth, B., et al. 2000. Differentiation of the epithelial apical junctional complex during mouse preimplantation development: a role for Rab13 in the early maturation of the tight junction. *Mech. Dev.* 97:93-104.
- Yamamoto, Y., et al. 2003. Distinct roles of Rab 3B and Rab 13 in the polarized transport of apical, basolateral and tight junctional membrane proteins to the plasma membrane. *Biochem. Biophys. Res. Commun.* 308: 270-275.
- Morimoto, S., et al. 2005. Rab13 mediates the continuous endocytic recycling of occludin to the cell surface. *J. Biol. Chem.* 280: 2220-2228.
- Marzesco, A.M. and Zahraoui, A. 2005. Assay of Rab13 in regulating epithelial tight junction assembly. *Methods Enzymol.* 403: 182-193.

CHROMOSOMAL LOCATION

Genetic locus: Rab13 (mouse) mapping to 3 F1.

PRODUCT

Rab 13 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rab 13 shRNA Plasmid (m): sc-152627-SH and Rab 13 shRNA (m) Lentiviral Particles: sc-152627-V as alternate gene silencing products.

For independent verification of Rab 13 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152627A, sc-152627B and sc-152627C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rab 13 siRNA (m) is recommended for the inhibition of Rab 13 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Rab 13 (8E8E2): sc-517224 is recommended as a control antibody for monitoring of Rab 13 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rab 13 gene expression knockdown using RT-PCR Primer: Rab 13 (m)-PR: sc-152627-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.