



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Rab 28 siRNA (m): sc-152632

## BACKGROUND

Ras-related GTPases are GTP-dependent switches that operate signaling pathways in cell growth, metabolism and organelle trafficking. Rab 28, also known as MGC41862, is a 221 amino acid protein that belongs to the small GTPase superfamily and the Rab family. Rab 28 is only distantly related to other Rab family members, sharing 31 to 33% amino acid identity with Rab 1A, Rab 6, Rab 11 and Rab 13. Rab 28 is expressed as two isoforms, designated S and L respectively, that are produced by alternative splicing events. Isoform S is detected in most tissues including cortex, liver, kidney, skeletal muscle, adipose tissue, testis and urothelium, where as isoform L is predominantly expressed in testis. Existing as two alternatively spliced isoforms, the Rab 28 gene is conserved in chimpanzee, canine, bovine, mouse, rat, chicken and *C. elegans*, and maps to human chromosome 4p15.33. Pseudogenes of the Rab 28 gene are found on chromosomes 9 and X.

## REFERENCES

1. Brauers, A., Schürmann, A., Massmann, S., Mühl-Zürbes, P., Becker, W., Kainulainen, H., Lie, C. and Joost, H.G. 1996. Alternative mRNA splicing of the novel GTPase Rab28 generates isoforms with different C-termini. *Eur. J. Biochem.* 237: 833-840.
2. Pereira-Leal, J.B. and Seabra, M.C. 2000. The mammalian Rab family of small GTPases: definition of family and subfamily sequence motifs suggests a mechanism for functional specificity in the Ras superfamily. *J. Mol. Biol.* 301: 1077-1087.
3. Stenmark, H. and Olkkonen, V.M. 2001. The Rab GTPase family. *Genome Biol.* 2: REVIEWS3007.
4. Maurer-Stroh, S., Koranda, M., Benetka, W., Schneider, G., Sirota, F.L. and Eisenhaber, F. 2007. Towards complete sets of farnesylated and geranylgeranylated proteins. *PLoS Comput. Biol.* 3: e66.
5. Lee, S.H., Baek, K. and Dominguez, R. 2008. Large nucleotide-dependent conformational change in Rab28. *FEBS Lett.* 582: 4107-4111.
6. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 612994. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: Rab28 (mouse) mapping to 5 B3.

## PRODUCT

Rab 28 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rab 28 shRNA Plasmid (m): sc-152632-SH and Rab 28 shRNA (m) Lentiviral Particles: sc-152632-V as alternate gene silencing products.

For independent verification of Rab 28 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152632A, sc-152632B and sc-152632C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Rab 28 siRNA (m) is recommended for the inhibition of Rab 28 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rab 28 gene expression knockdown using RT-PCR Primer: Rab 28 (m)-PR: sc-152632-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.