



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

rabphilin-3AL siRNA (m): sc-152672

BACKGROUND

Rabphilin-3AL (rabphilin-3A-like), also known as RPH3AL or NOC2, is a cytoplasmic Rab GTPase effector. It contains one FYVE-type zinc finger and one Rab-binding (RBD) domain, but unlike its related protein, rabphilin-3A, rabphilin-3AL does not contain any C2 domains. Rabphilin-3AL is expressed in a variety of tissues, with highest levels found in kidney, skeletal muscle, pancreas, liver, ovary, stomach, heart and thyroid. It is believed to play a role regulating calcium-dependent secretory vesicle exocytosis in endocrine and exocrine cells. Via its RBD domain, rabphilin-3AL is capable of binding Rab 27a and, through this interaction, rabphilin-3AL is recruited to dense-core vesicles. With lower affinity, rabphilin-3AL can also bind Rab 3 and Rab 8 with its RBD domain. Through an interaction with Rab 3, rabphilin-3AL can inhibit G protein signaling in endocrine pancreas and positively regulate Insulin secretion. Rabphilin-3AL knockout mice display accumulation of secretory granules and irregular shape in exocrine cells.

REFERENCES

1. Cheviet, S., Waselle, L. and Regazzi, R. 2004. NOC-king out exocrine and endocrine secretion. *Trends Cell Biol.* 14: 525-528.
2. Matsumoto, M., Miki, T., Shibasaki, T., Kawaguchi, M., Shinozaki, H., Nio, J., Saraya, A., Koseki, H., Miyazaki, M., Iwanaga, T. and Seino, S. 2004. NOC2 is essential in normal regulation of exocytosis in endocrine and exocrine cells. *Proc. Natl. Acad. Sci. USA* 101: 8313-8318.
3. Fukuda, M., Kanno, E. and Yamamoto, A. 2004. Rabphilin and NOC2 are recruited to dense-core vesicles through specific interaction with Rab 27a in PC12 cells. *J. Biol. Chem.* 279: 13065-13075.
4. Shibasaki, T. and Seino, S. 2005. Physical and functional interaction of NOC2/Rab 3 in exocytosis. *Meth. Enzymol.* 403: 408-419.
5. Fukuda, M. and Yamamoto, A. 2005. Assay of the Rab-binding specificity of rabphilin and NOC2: target molecules for Rab 27. *Meth. Enzymol.* 403: 469-481.
6. Imai, A., Yoshie, S., Nashida, T., Shimomura, H. and Fukuda, M. 2006. Functional involvement of NOC2, a Rab 27 effector, in rat parotid acinar cells. *Arch. Biochem. Biophys.* 455: 127-135.

CHROMOSOMAL LOCATION

Genetic locus: Rph3al (mouse) mapping to 11 B5.

PRODUCT

rabphilin-3AL siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see rabphilin-3AL shRNA Plasmid (m): sc-152672-SH and rabphilin-3AL shRNA (m) Lentiviral Particles: sc-152672-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

rabphilin-3AL siRNA (m) is recommended for the inhibition of rabphilin-3AL expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor rabphilin-3AL gene expression knockdown using RT-PCR Primer: rabphilin-3AL (m)-PR: sc-152672-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.