



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# RACK7 siRNA (m): sc-152674

## BACKGROUND

Members of the protein kinase C (PKC) family play a key regulatory role in a variety of cellular functions including cell growth and differentiation, gene expression, hormone secretion and membrane function. Receptor for activated C kinases, termed RACKs, are intracellular receptors for activated PKC that serve as anchors and may be involved in the activation-induced translocation of PKC. RACK7 (receptor for activated C kinase 7), also known as ZMYND8 (zinc finger MYND domain-containing protein 8), PRKCBP1 (protein kinase C (PKC)-binding protein 1) or PRO2893, is a widely expressed protein with predominant expression in pancreas, lung, placenta and brain. RACK7 contains one bromodomain, one PHD-type zinc finger, one MYND-type zinc finger and one PWWP domain. Via its C-terminus, RACK7 interacts with PKC  $\beta$  and is believed to play a role in PKC signaling and function as a transcription regulator. In response to DNA damage, RACK7 is phosphorylated by ATM or ATR. In addition, multiple isoforms exist for RACK7.

## REFERENCES

1. Fossey, S.C., et al. 2000. Identification and characterization of PRKCBP1, a candidate RACK-like protein. *Mamm. Genome* 11: 919-925.
2. Zhang, Y., et al. 2001. Identification of differentially expressed genes following treatment of monkey kidney cells with the mycotoxin fumonisin B<sub>1</sub>. *Food Chem. Toxicol.* 39: 45-53.
3. Park, J., et al. 2002. Mutation profiling of mismatch repair-deficient colorectal cancers using an in silico genome scan to identify coding microsatellites. *Cancer Res.* 62: 1284-1288.
4. Ansieau, S. and Sergeant, A. 2003. BS69 and RACK7, a potential novel class of tumor suppressor genes. *Pathol. Biol.* 51: 397-399.
5. Westendorf, J.J. and Koka, S. 2004. Identification of FHOD1-binding proteins and mechanisms of FHOD1-regulated Actin dynamics. *J. Cell. Biochem.* 92: 29-41.
6. Miles, R.R., et al. 2005. Analysis of BCL6-interacting proteins by tandem mass spectrometry. *Mol. Cell. Proteomics* 4: 1898-1909.

## CHROMOSOMAL LOCATION

Genetic locus: Prkcbp1 (mouse) mapping to 2 H3.

## PRODUCT

RACK7 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RACK7 shRNA Plasmid (m): sc-152674-SH and RACK7 shRNA (m) Lentiviral Particles: sc-152674-V as alternate gene silencing products.

For independent verification of RACK7 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152674A, sc-152674B and sc-152674C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

RACK7 siRNA (m) is recommended for the inhibition of RACK7 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RACK7 gene expression knockdown using RT-PCR Primer: RACK7 (m)-PR: sc-152674-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.