



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

retbindin siRNA (m): sc-152813

BACKGROUND

Retbindin, also known as RTBDN, is a 229 amino acid secreted protein that belongs to the folate receptor family. Expressed in retina, the gene that encodes retbindin exists as two alternatively spliced isoforms. Containing six exons and spanning approximately 5 kb, the retbindin gene is conserved in chimpanzee, canine, bovine, mouse and rat, and maps to human chromosome 19p13.13. Chromosome 19 consists of approximately 63 million bases and makes up over 2% of human genomic DNA. Chromosome 19 includes a diversity of interesting genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG families, and Fc α receptors. Key genes for eye color and hair color also map to chromosome 19.

REFERENCES

- Trettel, F., Mantuano, E., Calabresi, V., Veneziano, L., Olsen, A.S., Georgescu, A., Gordon, L., Sabbadini, G., Frontali, M. and Jodice, C. 2000. A fine physical map of the CACNA1A gene region on 19p13.1-p13.2 chromosome. *Gene* 241: 45-50.
- Wistow, G., Bernstein, S.L., Wyatt, M.K., Ray, S., Behal, A., Touchman, J.W., Bouffard, G., Smith, D. and Peterson, K. 2002. Expressed sequence tag analysis of human retina for the NEIBank Project: retbindin, an abundant, novel retinal cDNA and alternative splicing of other retina-preferred gene transcripts. *Mol. Vis.* 8: 196-204.
- Grimwood, J., Gordon, L.A., Olsen, A., Terry, A., Schmutz, J., Lamerdin, J., Hellsten, U., Goodstein, D., Couronne, O., Tran-Gyamfi, M., Aerts, A., Altherr, M., Ashworth, L., Bajorek, E., Black, S., Branscomb, E., et al. 2004. The DNA sequence and biology of human chromosome 19. *Nature* 428: 529-535.
- Online Mendelian Inheritance in Man, OMIM[™]. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 609553. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Kelemen, L.E. 2006. The role of folate receptor α in cancer development, progression and treatment: cause, consequence or innocent bystander? *Int. J. Cancer* 119: 243-250.
- Tian, X., Gautron, J., Monget, P. and Pascal, G. 2010. What makes an egg unique? Clues from evolutionary scenarios of egg-specific genes. *Biol. Reprod.* 83: 893-900.
- SWISS-PROT/TrEMBL (Q9BSG5). World Wide Web URL: <http://www.uniprot.org/uniprot/Q9BSG5>

CHROMOSOMAL LOCATION

Genetic locus: Rtdn (mouse) mapping to 8 C3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

retbindin siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see retbindin shRNA Plasmid (m): sc-152813-SH and retbindin shRNA (m) Lentiviral Particles: sc-152813-V as alternate gene silencing products.

For independent verification of retbindin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152813A and sc-152813B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

retbindin siRNA (m) is recommended for the inhibition of retbindin expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor retbindin gene expression knockdown using RT-PCR Primer: retbindin (m)-PR: sc-152813-PR (20 μ l). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.