



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rhotekin 2 siRNA (m): sc-152861

BACKGROUND

Rho, the Ras-related small GTPase, is responsible for the regulation of Actin-based cytoskeletal structures including stress fibers, focal adhesions and the contractile ring apparatus. Rho proteins function as molecular switches that are able to turn cytokinesis on and off. Although little is known about signaling downstream of Rho, a host of putative Rho effector proteins have been described, including rhotillin, Rhotekin, citron and the serine/threonine kinase, protein kinase N. Rhotekin 2, also known as PLEKHK1 (Pleckstrin homology domain-containing family K member 1), is a 609 amino acid protein that is expressed in bone marrow-derived cells, lymphocytes and CD4 positive T-cells. Due to this expression pattern, it is likely that Rhotekin 2 participates in lymphopoiesis. Rhotekin 2 contains one PH domain, a motif that is present in a variety of proteins that are involved in intracellular signaling. There are three isoforms of Rhotekin that are produced as a result of alternative splicing events.

REFERENCES

1. Reid, T., Furuyashiki, T., Ishizaki, T., Watanabe, G., Watanabe, N., Fujisawa, K., Morii, N., Madaule, P. and Narumiya, S. 1996. Rhotekin, a new putative target for Rho bearing homology to a serine/threonine kinase, PKN, and rhotillin in the Rho-binding domain. *J. Biol. Chem.* 271: 13556-13560.
2. Collier, F.M., Gregorio-King, C.C., Gough, T.J., Talbot, C.D., Walder, K. and Kirkland, M.A. 2004. Identification and characterization of a lymphocytic Rho-GTPase effector: Rhotekin-2. *Biochem. Biophys. Res. Commun.* 324: 1360-1369.
3. Chen, D., Zhao, M. and Mundy, G.R. 2004. Bone morphogenetic proteins. *Growth Factors* 22: 233-241.
4. Gregorio-King, C.C., Gough, T., Van Der Meer, G.J., Hosking, J.B., Waugh, C.M., McLeod, J.L., Collier, F.M. and Kirkland, M.A. 2004. Mechanisms of resistance to the cytotoxic effects of oxysterols in human leukemic cells. *J. Steroid Biochem. Mol. Biol.* 88: 311-320.
5. Ito, H., Iwamoto, I., Mizutani, K., Morishita, R., Deguchi, T., Nozawa, Y., Asano, T. and Nagata, K. 2006. Possible interaction of a Rho effector, Rhotekin, with a PDZ-protein, PIST, at synapses of hippocampal neurons. *Neurosci. Res.* 56: 165-171.
6. Williams, D.A., Zheng, Y. and Cancelas, J.A. 2008. Rho GTPases and regulation of hematopoietic stem cell localization. *Meth. Enzymol.* 439: 365-393.
7. Nagata, K., Ito, H., Iwamoto, I., Morishita, R. and Asano, T. 2009. Interaction of a multi-domain adaptor protein, vinexin, with a Rho-effector, Rhotekin. *Med. Mol. Morphol.* 42: 9-15.

CHROMOSOMAL LOCATION

Genetic locus: Rtnk2 (mouse) mapping to 10 B5.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Rhotekin 2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rhotekin 2 shRNA Plasmid (m): sc-152861-SH and Rhotekin 2 shRNA (m) Lentiviral Particles: sc-152861-V as alternate gene silencing products.

For independent verification of Rhotekin 2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152861A, sc-152861B and sc-152861C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rhotekin 2 siRNA (m) is recommended for the inhibition of Rhotekin 2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rhotekin 2 gene expression knockdown using RT-PCR Primer: Rhotekin 2 (m)-PR: sc-152861-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.