



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Ribosomal Protein L17 siRNA (m): sc-152898

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein L17, also known as RPL17, rPL23, PD-1 or 60S Ribosomal protein L17, is a 184 amino acid protein that is expressed in pancreas, lung, colon, cystic duct, gall bladder, kidney and liver and belongs to the ribosomal protein L22P family. Localized to the cytoplasm, it has been suggested that Ribosomal protein L17 may influence sexual differentiation of Area X and RA, potentially regulating the genesis and/or survival of neurons of juvenile zebra finches.

REFERENCES

1. Wool, I.G., Chan, Y.L. and Glück, A. 1995. Structure and evolution of mammalian ribosomal proteins. *Biochem. Cell Biol.* 73: 933-947.
2. Frigerio, J.M., Dagorn, J.C. and Iovanna, J.L. 1995. Cloning, sequencing and expression of the L5, L21, L27a, L28, S5, S9, S10 and S29 human ribosomal protein mRNAs. *Biochim. Biophys. Acta* 1262: 64-68.
3. Kenmochi, N., Kawaguchi, T., Rozen, S., Davis, E., Goodman, N., Hudson, T.J., Tanaka, T. and Page, D.C. 1998. A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509-523.
4. Kusuda, J., Hirai, M., Tanuma, R., Hirata, M. and Hashimoto, K. 1999. Genomic structure and chromosome location of RPL27A/Rpl27a, the genes encoding human and mouse ribosomal protein L27A. *Cytogenet. Cell Genet.* 85: 248-251.
5. Bortoluzzi, S., d'Alessi, F., Romualdi, C. and Danieli, G.A. 2001. Differential expression of genes coding for ribosomal proteins in different human tissues. *Bioinformatics* 17: 1152-1157

CHROMOSOMAL LOCATION

Genetic locus: Rpl17 (mouse) mapping to 18 E3.

PRODUCT

Ribosomal Protein L17 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribosomal Protein L17 shRNA Plasmid (m): sc-152898-SH and Ribosomal Protein L17 shRNA (m) Lentiviral Particles: sc-152898-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ribosomal Protein L17 siRNA (m) is recommended for the inhibition of Ribosomal Protein L17 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Ribosomal Protein L17 (C-8): sc-515904 is recommended as a control antibody for monitoring of Ribosomal Protein L17 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ribosomal Protein L17 gene expression knockdown using RT-PCR Primer: Ribosomal Protein L17 (m)-PR: sc-152898-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.