



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

RIC-8B siRNA (m): sc-152955

BACKGROUND

The Ras superfamily of GTPases can be subdivided into the Ras, Rho/Rac, Sar, Rab, Arf, Rap and Ran subfamilies, all of which control multiple aspects of cell function, including cytoskeletal rearrangement, nuclear signaling and cell growth. The Ras superfamily of GTPases function as regulated switches that toggle between a biologically active GTP-bound and an inactive GDP-bound form. This activation is catalyzed by guanine nucleotide exchange factors (GEFs). RIC-8B (resistance to inhibitors of cholinesterase 8 homolog B), also known as Synembryn-B (brain synembryn) or hSyn in human, is a 536 amino acid cytoplasmic protein that can activate several G_{α} proteins, including $G_{\alpha i-1}$, $G_{\alpha q}$ and $G_{\alpha o}$. Functioning as a guanine nucleotide exchange factor, RIC-8B binds to GDP-associated substrates and exchanges bound GDP for free GTP. RIC-8B has been localized to mature olfactory sensory neurons as well as in a few regions in the brain and has been shown to play a role in olfactory receptor expression and signaling. RIC-8B is expressed as multiple isoforms due to alternative splicing events.

REFERENCES

1. Miller, K.G., et al. 2000. RIC-8 (synembryn): a novel conserved protein that is required for $G_{\alpha q}$ signaling in the *C. elegans* nervous system. *Neuron* 27: 289-299.
2. Klattenhoff, C., et al. 2003. Human brain synembryn interacts with $G_{\alpha s}$ and $G_{\alpha q}$ and is translocated to the plasma membrane in response to isoproterenol and carbachol. *J. Cell. Physiol.* 195: 151-157.
3. Malik, S., et al. 2005. RIC-8 enhances G protein $\beta\gamma$ -dependent signaling in response to $\beta\gamma$ -binding peptides in intact cells. *Mol. Pharmacol.* 68: 129-136.
4. Charlie, N.K., et al. 2006. The dunce cAMP phosphodiesterase PDE-4 negatively regulates $G_{\alpha s}$ -dependent and $G_{\alpha s}$ -independent cAMP pools in the *Caenorhabditis elegans* synaptic signaling network. *Genetics* 173: 111-130.

CHROMOSOMAL LOCATION

Genetic locus: Ric8b (mouse) mapping to 10 C1.

PRODUCT

RIC-8B siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RIC-8B shRNA Plasmid (m): sc-152955-SH and RIC-8B shRNA (m) Lentiviral Particles: sc-152955-V as alternate gene silencing products.

For independent verification of RIC-8B (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152955A, sc-152955B and sc-152955C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RIC-8B siRNA (m) is recommended for the inhibition of RIC-8B expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RIC-8B gene expression knockdown using RT-PCR Primer: RIC-8B (m)-PR: sc-152955-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.