



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rim2 siRNA (m): sc-152965

BACKGROUND

Rab 3, a neural/neuroendocrine-specific member of the Rab family, is involved in Ca^{2+} -regulated exocytosis. Rab 3 functions in an inhibitory capacity by controlling the recruitment of secretory vesicles into a releasable pool at the plasma membrane. Rim (Rab 3 interacting molecule), a putative effector protein for Rab3s, is composed of an N-terminal zinc-finger motif and C-terminal PDZ and C2 domains. Rim exists as two variants, Rim1 and Rim2, produced by alternative splicing. The 3'-end of the Rim2 gene produces an independent mRNA that encodes a smaller protein referred to as Nim2, which like Rim, also regulates exocytosis. Rim serves as a Rab 3-dependent regulator of synaptic-vesicle fusion by forming a GTP-dependent complex between synaptic plasma membranes and docked synaptic vesicles. Both Rim1 and Rim2 can bind to cAMP-GEFII, which is a direct target of cAMP in regulated exocytosis and is responsible for cAMP-dependent, PKA-dependent exocytosis. Rim also localizes on the plasma membrane of INS-1E cells and pancreatic β -cells. Rab 3 binding domain of Rim enhances glucose-stimulated secretion in intact cells and Ca^{2+} -stimulated exocytosis in permeabilized cells, suggesting that Rim may also play a regulatory role in Insulin secretion.

REFERENCES

- Wang, Y., Okamoto, M., Schmitz, F., Hofmann, K. and Sudhof, T.C. 1997. Rim is a putative Rab 3 effector in regulating synaptic-vesicle fusion. *Nature* 388: 593-598.
- Coppola, T., Perret-Menoud, V., Luthi, S., Farnsworth, C.C., Glomset, J.A. and Regazzi, R. 1999. Disruption of Rab 3-calmodulin interaction, but not other effector interactions, prevents Rab 3 inhibition of exocytosis. *EMBO J.* 18: 5885-5891.
- Ozaki, N., Shibasaki, T., Kashima, Y., Miki, T., Takahashi, K., Ueno, H., Sunaga, Y., Yano, H., Matsuura, Y., Iwanaga, T., Takai, Y. and Seino, S. 2000. cAMP-GEFII is a target of cAMP in regulated exocytosis. *Nat. Cell Biol.* 2: 805-811.
- Wang, Y., Sugita, S. and Sudhof, T.C. 2000. The Rim/Nim family of neuronal C2 domain proteins. Interactions with Rab 3 and a new class of Src homology 3 domain proteins. *J. Biol. Chem.* 275: 20043-20044.
- Iezzi, M., Regazzi, R. and Wollheim, C.B. 2000. The Rab3-interacting molecule Rim is expressed in pancreatic β -cells and is implicated in Insulin exocytosis. *FEBS Lett.* 474: 66-70.
- Haynes, L.P., Evans, G.J., Morgan, A. and Burgoyne, R.D. 2001. A direct inhibitory role for the Rab 3-specific effector, Noc2, in Ca^{2+} -regulated exocytosis in neuroendocrine cells. *J. Biol. Chem.* 276: 9726-9732.

CHROMOSOMAL LOCATION

Genetic locus: Rims2 (mouse) mapping to 15 B3.1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Rim2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rim2 shRNA Plasmid (m): sc-152965-SH and Rim2 shRNA (m) Lentiviral Particles: sc-152965-V as alternate gene silencing products.

For independent verification of Rim2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152965A, sc-152965B and sc-152965C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rim2 siRNA (m) is recommended for the inhibition of Rim2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Rim2 (63-M7): sc-100842 is recommended as a control antibody for monitoring of Rim2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rim2 gene expression knockdown using RT-PCR Primer: Rim2 (m)-PR: sc-152965-PR (20 μ l). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.