



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

RN-tre siRNA (m): sc-152987

BACKGROUND

GTPase-activating proteins (GAPs) accelerate the intrinsic rate of GTP hydrolysis of Ras-related proteins, resulting in downregulation of their active form. RN-tre (related to the N-terminus of tre), also known as USP6NL (USP6 N-terminal like) or TRE2NL, is an 828 amino acid protein that contains one Rab-GAP TBC domain. Expressed in a wide variety of tissues, RN-tre functions as a GTPase-activating protein for Rab 5A and is thought to be involved in receptor trafficking, as well as in the inhibition of EGFR internalization. In response to DNA damage, RN-tre may be phosphorylated by ATM or ATR. The gene encoding RN-tre maps to human chromosome 10, which houses over 1,200 genes and comprises nearly 4.5% of the human genome.

REFERENCES

1. Nomura, N., et al. 1994. Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1. *DNA Res.* 1: 27-35.
2. Matůšková, B., et al. 1996. RN-tre identifies a family of tre-related proteins displaying a novel potential protein binding domain. *Oncogene* 12: 2563-2571.
3. Matůšková, B., et al. 1996. RN-tre specifically binds to the SH3 domain of eps8 with high affinity and confers growth advantage to NIH3T3 upon carboxy-terminal truncation. *Oncogene* 12: 2679-2688.
4. Lanzetti, L., et al. 2000. The Eps8 protein coordinates EGF receptor signalling through Rac and trafficking through Rab5. *Nature* 408: 374-377.
5. Martinu, L., et al. 2002. Endocytosis of epidermal growth factor receptor regulated by Grb2-mediated recruitment of the Rab5 GTPase-activating protein RN-tre. *J. Biol. Chem.* 277: 50996-51002.
6. Lanzetti, L., et al. 2004. Rab5 is a signalling GTPase involved in Actin remodelling by receptor tyrosine kinases. *Nature* 429: 309-314.
7. Lanzetti, L., et al. 2007. Regulation of the Rab5 GTPase-activating protein RN-tre by the dual specificity phosphatase Cdc14A in human cells. *J. Biol. Chem.* 282: 15258-15270.

CHROMOSOMAL LOCATION

Genetic locus: *Usp6nl* (mouse) mapping to 2 A1.

PRODUCT

RN-tre siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RN-tre shRNA Plasmid (m): sc-152987-SH and RN-tre shRNA (m) Lentiviral Particles: sc-152987-V as alternate gene silencing products.

For independent verification of RN-tre (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152987A, sc-152987B and sc-152987C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RN-tre siRNA (m) is recommended for the inhibition of RN-tre expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RN-tre gene expression knockdown using RT-PCR Primer: RN-tre (m)-PR: sc-152987-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.