



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ROM1 siRNA (m): sc-153060

BACKGROUND

ROM1 (retinal outer segment membrane protein 1), also known as ROM, RP7, ROSP1 or TSPAN23, is a 351 amino acid integral membrane protein found in the photoreceptor disk rim of the eye. Belonging to the PRPH2/ROM1 family, ROM1 exists as a homodimer but can also form heterodimers with retinal degeneration slow (RDS). ROM1 is essential for disk morphogenesis and may also function as an adhesion molecule involved in the stabilization and compaction of outer segment disks or in the maintenance of the curvature of the rim. Certain defects in the gene encoding ROM1 have been associated with retinitis pigmentosa, a group of genetic eye conditions characterized by the progressive loss of photoreceptor cells that may potentially lead to blindness.

REFERENCES

1. Bascom, R.A., Liu, L., Humphries, P., Fishman, G.A., Murray, J.C. and McInnes, R.R. 1993. Polymorphisms and rare sequence variants at the ROM1 locus. *Hum. Mol. Genet.* 2: 1975-1977.
2. Taylor, B.A., Phillips, S.J., Bascom, R.A. and McInnes, R.R. 1994. The retinal outer segment membrane protein-1 gene (ROM1) maps to the proximal end of mouse chromosome 19. *Genomics* 23: 510-511.
3. Bascom, R.A., Liu, L., Heckenlively, J.R., Stone, E.M. and McInnes, R.R. 1995. Mutation analysis of the ROM1 gene in retinitis pigmentosa. *Hum. Mol. Genet.* 4: 1895-1902.
4. Gould, D.J., Petersen-Jones, S.M., Lin, C.T. and Sargan, D.R. 1997. Cloning of canine rom-1 and its investigation as a candidate gene for generalized progressive retinal atrophies in dogs. *Anim. Genet.* 28: 391-396.
5. Dryja, T.P., Hahn, L.B., Kajiwar, K. and Berson, E.L. 1997. Dominant and digenic mutations in the peripherin/RDS and ROM1 genes in retinitis pigmentosa. *Invest. Ophthalmol. Vis. Sci.* 38: 1972-1982.
6. Runte, M., Dekomien, G. and Epplen, J.T. 2000. Evaluation of RDS/Peripherin and ROM1 as candidate genes in generalised progressive retinal atrophy and exclusion of digenic inheritance. *Anim. Genet.* 31: 223-227.
7. Boesze-Battaglia, K., Dispoto, J. and Kahoe, M.A. 2002. Association of a photoreceptor-specific tetraspanin protein, ROM1, with triton X-100-resistant membrane rafts from rod outer segment disk membranes. *J. Biol. Chem.* 277: 41843-41849.
8. Hartong, D.T., Berson, E.L. and Dryja, T.P. 2006. Retinitis pigmentosa. *Lancet* 368: 1795-1809.
9. Ivings, L., Towns, K.V., Matin, M.A., Taylor, C., Ponchel, F., Grainger, R.J., Ramesar, R.S., Mackey, D.A. and Inglehearn, C.F. 2008. Evaluation of splicing efficiency in lymphoblastoid cell lines from patients with splicing-factor retinitis pigmentosa. *Mol. Vis.* 14: 2357-2366.

CHROMOSOMAL LOCATION

Genetic locus: Rom1 (mouse) mapping to 19 A.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

ROM1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ROM1 shRNA Plasmid (m): sc-153060-SH and ROM1 shRNA (m) Lentiviral Particles: sc-153060-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ROM1 siRNA (m) is recommended for the inhibition of ROM1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ROM1 gene expression knockdown using RT-PCR Primer: ROM1 (m)-PR: sc-153060-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.