



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TCEAL7 siRNA (m): sc-153069

BACKGROUND

TCEAL7 (transcription elongation factor A protein-like 7), also known as transcription elongation factor S-II protein-like 7, is a 100 amino acid nuclear protein that belongs to the TFS-II family and the TFA subfamily. While highly expressed in normal and fetal brain tissues, TCEAL7 is weakly expressed in uterus and ovary. In addition, TCEAL7 is down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. TCEAL7 plays a role in the negative regulation of NF κ B signaling at the basal level by modulating transcriptional activity of NF κ B on its target gene promoters. Associating with cyclin D1 promoter containing Myc E-box sequence, TCEAL7 transcriptionally represses cyclin D1 expression. Acting in both ALT (alternative lengthening of telomeres) and telomerase-positive cell lines, TCEAL7 regulates telomerase reverse transcriptase expression and telomerase activity.

REFERENCES

- Chien, J., Staub, J., Avula, R., Zhang, H., Liu, W., Hartmann, L.C., Kaufmann, S.H., Smith, D.I. and Shridhar, V. 2005. Epigenetic silencing of TCEAL7 (Bex4) in ovarian cancer. *Oncogene* 24: 5089-5100.
- Chien, J., Narita, K., Rattan, R., Giri, S., Shridhar, R., Staub, J., Belefard, D., Lai, J., Roberts, L.R., Molina, J., Kaufmann, S.H., Prendergast, G.C. and Shridhar, V. 2008. A role for candidate tumor-suppressor gene TCEAL7 in the regulation of c-Myc activity, cyclin D1 levels and cellular transformation. *Oncogene* 27: 7223-7234.
- Peedicayil, A., Vierkant, R.A., Shridhar, V., Schildkraut, J.M., Armasu, S., Hartmann, L.C., Fridley, B.L., Cunningham, J.M., Phelan, C.M., Sellers, T.A. and Goode, E.L. 2009. Polymorphisms in TCEAL7 and risk of epithelial ovarian cancer. *Gynecol. Oncol.* 114: 260-264.
- Reams, R.R., Agrawal, D., Davis, M.B., Yoder, S., Odedina, F.T., Kumar, N., Higginbotham, J.M., Akinremi, T., Suther, S. and Soliman, K.F. 2009. Microarray comparison of prostate tumor gene expression in African-American and Caucasian American males: a pilot project study. *Infect. Agents Cancer* 4: S3.
- Shi, X. and Garry, D.J. 2010. Myogenic regulatory factors transactivate the TCEAL7 gene and modulate muscle differentiation. *Biochem. J.* 428: 213-221.
- Lafferty-Whyte, K., Bilsland, A., Hoare, S.F., Burns, S., Zaffaroni, N., Cairney, C.J. and Keith, W.N. 2010. TCEAL7 inhibition of c-Myc activity in alternative lengthening of telomeres regulates hTERT expression. *Neoplasia* 12: 405-414.
- Rattan, R., Narita, K., Chien, J., Maguire, J.L., Shridhar, R., Giri, S. and Shridhar, V. 2010. TCEAL7, a putative tumor suppressor gene, negatively regulates NF κ B pathway. *Oncogene* 29: 1362-1373.

CHROMOSOMAL LOCATION

Genetic locus: Tceal7 (mouse) mapping to X F1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TCEAL7 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCEAL7 shRNA Plasmid (m): sc-153069-SH and TCEAL7 shRNA (m) Lentiviral Particles: sc-153069-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TCEAL7 siRNA (m) is recommended for the inhibition of TCEAL7 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCEAL7 gene expression knockdown using RT-PCR Primer: TCEAL7 (m)-PR: sc-153069-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.