



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# RRP22 siRNA (m): sc-153132

## BACKGROUND

Members of the Ras superfamily of small GTP-binding proteins are critical mediators of diverse cell signaling pathways, including those leading to cell proliferation, cytoskeletal organization and secretion. The counter-conversion of the active GTP-bound form of these proteins to their inactive GDP-bound form is influenced by two types of regulatory proteins: those that alter the intrinsic GTPase activity of the GTP-binding proteins and those that alter the rate of GDP/GTP exchange. Guanine nucleotide-releasing factors (GRFs) increase the GDP dissociation rate, while GDP-dissociation inhibitors (GDIs) decrease the dissociation rate. RRP22 (Ras-related protein on chromosome 22), also known as RASL10A, is a 203 amino acid cell membrane protein belonging to the RAS family of the small GTPase superfamily. Unlike Ras, RRP22 inhibits cell growth and promotes caspase-independent cell death. Existing as two alternatively spliced isoforms, RRP22 is considered a potential tumor suppressor.

## REFERENCES

- Hall, A. 1990. The cellular functions of small GTP-binding proteins. *Science* 249: 635-640.
- Bourne, H.R., Sanders, D.A. and McCormick, F. 1990. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348: 125-132.
- Scheffzek, K., Klebe, C., Fritz-Wolf, K., Kabsch, W. and Wittinghofer, A. 1995. Crystal structure of the nuclear Ras-related protein Ran in its GDP-bound form. *Nature* 374: 378-381.
- Zucman-Rossi, J., Legoix, P. and Thomas, G. 1996. Identification of new members of the Gas2 and Ras families in the 22q12 chromosome region. *Genomics* 38: 247-254.
- Roa, M., Paumet, F., Le Mao, J., David, B. and Blank, U. 1997. Involvement of the Ras-like GTPase rab3d in RBL-2H3 mast cell exocytosis following stimulation via high affinity IgE receptors (Fc epsilonRI). *J. Immunol.* 159: 2815-2823.
- Elam, C., Hesson, L., Vos, M.D., Eckfeld, K., Ellis, C.A., Bell, A., Krex, D., Birrer, M.J., Latif, F. and Clark, G.J. 2005. RRP22 is a farnesylated, nucleolar, Ras-related protein with tumor suppressor potential. *Cancer Res.* 65: 3117-3125.
- Sayagues, J.M., Taberner, M.D. and Maíllo, A. 2007. Cytogenetic alterations in meningioma tumors and their impact on disease outcome. *Med. Clin.* 128: 226-232.

## CHROMOSOMAL LOCATION

Genetic locus: Rasl10a (mouse) mapping to 11 A1.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

RRP22 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RRP22 shRNA Plasmid (m): sc-153132-SH and RRP22 shRNA (m) Lentiviral Particles: sc-153132-V as alternate gene silencing products.

For independent verification of RRP22 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153132A and sc-153132B.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

RRP22 siRNA (m) is recommended for the inhibition of RRP22 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RRP22 gene expression knockdown using RT-PCR Primer: RRP22 (m)-PR: sc-153132-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.