



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rslcan-8 siRNA (m): sc-153157

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. Rslcan-8 (regulator of sex-limitation candidate 8), whose alternative names include zinc finger protein 759 (ZFP759) or BC028265, is a 721 amino acid KRAB zinc-finger protein belonging to the regulation of sex-limitation (Rsl) family, which is involved in the regulation of sex-dependent gene expression. Members of the Rsl family act on a variety of male-specific liver genes including a cytochrome P450 known as Cyp 2d-9 and mouse urinary proteins (MUPs). The gene encoding Rslcan-8 maps to mouse chromosome 13 B3.

REFERENCES

- Jiang, P.P., Frederick, K., Hansen, T.H. and Miller, R.D. 1996. Localization of the mouse gene releasing sex-limited expression of Slp. Proc. Natl. Acad. Sci. USA 93: 913-917.
- Tullis, K.M., Krebs, C.J., Leung, J.Y. and Robins, D.M. 2003. The regulator of sex-limitation gene, Rsl, enforces male-specific liver gene expression by negative regulation. Endocrinology 144: 1854-1860.
- Krebs, C.J., Larkins, L.K., Price, R., Tullis, K.M., Miller, R.D. and Robins, D.M. 2003. Regulator of sex-limitation (Rsl) encodes a pair of KRAB zinc-finger genes that control sexually dimorphic liver gene expression. Genes Dev. 17: 2664-2674.
- Urrutia, R. 2003. KRAB-containing zinc-finger repressor proteins. Genome Biol. 4: 231.
- Krebs, C.J., Larkins, L.K., Khan, S.M. and Robins, D.M. 2005. Expansion and diversification of KRAB zinc-finger genes within a cluster including regulator of sex-limitation 1 and 2. Genomics 85: 752-761.
- Itokawa, Y., Yanagawa, T., Yamakawa, H., Watanabe, N., Koga, H. and Nagase, T. 2009. KAP1-independent transcriptional repression of SCAN-KRAB-containing zinc finger proteins. Biochem. Biophys. Res. Commun. 388: 689-694.
- Krebs, C.J., Khan, S., MacDonald, J.W., Sorenson, M. and Robins, D.M. 2009. Regulator of sex-limitation KRAB zinc finger proteins modulate sex-dependent and -independent liver metabolism. Physiol. Genomics 38: 16-28.

CHROMOSOMAL LOCATION

Genetic locus: Zfp759 (mouse) mapping to 13 B3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Rslcan-8 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rslcan-8 shRNA Plasmid (m): sc-153157-SH and Rslcan-8 shRNA (m) Lentiviral Particles: sc-153157-V as alternate gene silencing products.

For independent verification of Rslcan-8 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153157A, sc-153157B and sc-153157C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rslcan-8 siRNA (m) is recommended for the inhibition of Rslcan-8 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rslcan-8 gene expression knockdown using RT-PCR Primer: Rslcan-8 (m)-PR: sc-153157-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.