



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



RWDD2B siRNA (m): sc-153182

BACKGROUND

RWDD2B (RWD domain containing protein 2B), also known as C21orf6, is a 319 amino acid protein that is ubiquitously expressed. RWDD2B contains one RWD domain, a conserved region of about 110 amino acid residues. RWD domains are found in many RING finger proteins, DEAD-like helicases and WD repeat containing proteins. It is believed that RWD domains may be involved in protein interaction. The gene that encodes RWDD2B maps to chromosome 21, which makes up about 1.5% of the human genome. Down syndrome, also known as trisomy 21, is the disease most commonly associated with chromosome 21. Alzheimer's disease, Jervell and Lange-Nielsen syndrome and amyotrophic lateral sclerosis are also associated with chromosome 21. Translocations are found to occur between chromosome 21 and 8, and chromosome 21 and 12, in certain leukemias.

REFERENCES

1. Tyson, J., Tranebjaerg, L., Bellman, S., Wren, C., Taylor, J.F., Bathen, J., Aslaksen, B., Sorland, S.J., Lund, O., Malcolm, S., Pembrey, M., Bhattacharya, S. and Bitner-Glindzicz, M. 1997. IsK and KvLQT1: mutation in either of the two subunits of the slow component of the delayed rectifier potassium channel can cause Jervell and Lange-Nielsen syndrome. *Hum. Mol. Genet.* 6: 2179-2185.
2. Müller, S., Stanyon, R., Finelli, P., Archidiacono, N. and Wienberg, J. 2000. Molecular cytogenetic dissection of human chromosomes 3 and 21 evolution. *Proc. Natl. Acad. Sci. USA* 97: 206-211.
3. Orti, R., Rachidi, M., Vialard, F., Toyama, K., Lopes, C., Taudien, S., Rosenthal, A., Yspo, M.L., Sinet, P.M. and Delabar, J.M. 2000. Characterization of a novel gene, C21orf6, mapping to a critical region of chromosome 21q22.1 involved in the monosomy 21 phenotype and of its murine ortholog, orf5. *Genomics* 64: 203-210.
4. Doerks, T., Copley, R.R., Schultz, J., Ponting, C.P. and Bork, P. 2002. Systematic identification of novel protein domain families associated with nuclear functions. *Genome Res.* 12: 47-56.
5. Mao, R., Wang, X., Spitznagel, E.L., Jr., Frelin, L.P., Ting, J.C., Ding, H., Kim, J.W., Ruczinski, I., Downey, T.J. and Pevsner, J. 2005. Primary and secondary transcriptional effects in the developing human Down syndrome brain and heart. *Genome Biol.* 6: R107.
6. Robakis, N.K. 2006. The discovery and mapping to chromosome 21 of the Alzheimer's amyloid gene: history revised. *J. Alzheimers Dis.* 10: 453-455.
7. Aït Yahya-Graison, E., Aubert, J., Dauphinot, L., Rivals, I., Prieur, M., Golfier, G., Rossier, J., Personnaz, L., Creau, N., Blehaut, H., Robin, S., Delabar, J.M. and Potier, M.C. 2007. Classification of human chromosome 21 gene-expression variations in Down syndrome: impact on disease phenotypes. *Am. J. Hum. Genet.* 81: 475-491.
8. Peterson, L.F., Boyapati, A., Ahn, E.Y., Biggs, J.R., Okumura, A.J., Lo, M.C., Yan, M. and Zhang, D.E. 2007. Acute myeloid leukemia with the 8q22;21q22 translocation: secondary mutational events and alternative t(8;21) transcripts. *Blood* 110: 799-805.

CHROMOSOMAL LOCATION

Genetic locus: Rwd2b (mouse) mapping to 16 C3.3.

PRODUCT

RWDD2B siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RWDD2B shRNA Plasmid (m): sc-153182-SH and RWDD2B shRNA (m) Lentiviral Particles: sc-153182-V as alternate gene silencing products.

For independent verification of RWDD2B (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153182A, sc-153182B and sc-153182C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RWDD2B siRNA (m) is recommended for the inhibition of RWDD2B expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RWDD2B gene expression knockdown using RT-PCR Primer: RWDD2B (m)-PR: sc-153182-PR (20 µl). Annealing temperature for the primers should be 55-60°C and the extension temperature should be 68-72°C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.