



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# SAMD1 siRNA (m): sc-153201

## BACKGROUND

SAMD1 (sterile  $\alpha$  motif domain containing 1), also known as atherin, is a 538 amino acid secreted protein that contains one SAM (sterile  $\alpha$  motif) domain. Amino acid sequences in SAMD1 shows extensive sequence conservation of the full-length sequences for human and rabbit SAMD1. SAMD1 is present only in atherosclerotic lesions, not in normal intima. Within lesions, SAMD1 is found both in the extracellular compartment and within foam cells. SAMD1 may play a role in atherogenesis by immobilizing LDL in the arterial wall. The SAMD1 gene maps to human chromosome 19p13.12. Chromosome 19 consists of approximately 63 million bases and makes up over 2% of human genomic DNA. Chromosome 19 includes a diversity of interesting genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family, and Fc $\alpha$  receptors. Key genes for eye color and hair color also map to chromosome 19.

## REFERENCES

- Trettel, F., Mantuano, E., Calabresi, V., Veneziano, L., Olsen, A.S., Georgescu, A., Gordon, L., Sabbadini, G., Frontali, M. and Jodice, C. 2000. A fine physical map of the CACNA1A gene region on 19p13.1-p13.2 chromosome. *Gene* 241: 45-50.
- Grimwood, J., Gordon, L.A., Olsen, A., Terry, A., Schmutz, J., Lamerdin, J., Hellsten, U., Goodstein, D., Couronne, O., Tran-Gyamfi, M., Aerts, A., Altherr, M., Ashworth, L., Bajorek, E., Black, S., Branscomb, E., et al. 2004. The DNA sequence and biology of human chromosome 19. *Nature* 428: 529-535.
- Lees, A.M., Deconinck, A.E., Campbell, B.D. and Lees, R.S. 2005. Atherin: a newly identified, lesion-specific, LDL-binding protein in human atherosclerosis. *Atherosclerosis* 182: 219-230.
- Baptista, J., Prigmore, E., Gribble, S.M., Jacobs, P.A., Carter, N.P. and Crolla, J.A. 2005. Molecular cytogenetic analyses of breakpoints in apparently balanced reciprocal translocations carried by phenotypically normal individuals. *Eur. J. Hum. Genet.* 13: 1205-1212.
- Lopez-Coviella, I., Mellott, T.M., Kovacheva, V.P., Berse, B., Slack, B.E., Zemelko, V., Schnitzler, A. and Blusztajn, J.K. 2006. Developmental pattern of expression of BMP receptors and Smads and activation of SAMD1 and SAMD5 by BMP9 in mouse basal forebrain. *Brain Res.* 1088: 49-56.
- Jensen, D.R., Martin, D.M., Gebarski, S., Sahoo, T., Brundage, E.K., Chinault, A.C., Otto, E.A., Chaki, M., Hildebrandt, F., Cheung, S.W. and Lesperance, M.M. 2009. A novel chromosome 19p13.12 deletion in a child with multiple congenital anomalies. *Am. J. Med. Genet. A* 149A: 396-402.

## CHROMOSOMAL LOCATION

Genetic locus: Samd1 (mouse) mapping to 8 C3.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

SAMD1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SAMD1 shRNA Plasmid (m): sc-153201-SH and SAMD1 shRNA (m) Lentiviral Particles: sc-153201-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

SAMD1 siRNA (m) is recommended for the inhibition of SAMD1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SAMD1 gene expression knockdown using RT-PCR Primer: SAMD1 (m)-PR: sc-153201-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.