



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SAT-1 siRNA (m): sc-153230

BACKGROUND

SAT-1 (sulfate anion transporter 1), also known as SLC26A1 (solute carrier family 26 (sulfate transporter), member 1) or EDM4, is a 701 amino acid multi-pass membrane protein that belongs to the SLC26A/SuIP transporter family of proteins. Members of this family are sulfate/anion transporter proteins that are well conserved in their genomic (number and size of exons) and protein (amino acid length among species) structures, yet they exhibit very restricted and distinct tissue expression patterns. SAT-1 is predominantly expressed in kidney and liver but can also be found at lower levels in spleen, small intestine, brain, pancreas, leukocytes, prostate, thymus, testis and colon. Localized to the plasma membrane, SAT-1 contains one STAS domain, 12 transmembrane domains, 2 N-glycosylation sites and multiple phosphorylation sites. Accepting oxalate as a cosubstrate, SAT-1 participates in transtubular sulfate reabsorption by mediating the exit of sulfate across the basolateral membrane.

REFERENCES

- Lohi, H., et al. 2000. Mapping of five new putative anion transporter genes in human and characterization of SLC26A6, a candidate gene for pancreatic anion exchanger. *Genomics* 70: 102-112.
- Vincourt, J.B., et al. 2002. Molecular cloning of SLC26A7, a novel member of the SLC26 sulfate/anion transporter family, from high endothelial venules and kidney. *Genomics* 79: 249-256.
- Lee, A., et al. 2003. The mouse sulfate anion transporter gene *Sat1* (*Slc26a1*): cloning, tissue distribution, gene structure, functional characterization, and transcriptional regulation thyroid hormone. *DNA Cell Biol.* 22: 19-31.
- Regeer, R.R., et al. 2003. Characterization of the human sulfate anion transporter (hSAT-1) protein and gene (SAT1; SLC26A1). *DNA Cell Biol.* 22: 107-117.

CHROMOSOMAL LOCATION

Genetic locus: *Slc26a1* (mouse) mapping to 5 F.

PRODUCT

SAT-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SAT-1 shRNA Plasmid (m): sc-153230-SH and SAT-1 shRNA (m) Lentiviral Particles: sc-153230-V as alternate gene silencing products.

For independent verification of SAT-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153230A, sc-153230B and sc-153230C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SAT-1 siRNA (m) is recommended for the inhibition of SAT-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SAT-1 gene expression knockdown using RT-PCR Primer: SAT-1 (m)-PR: sc-153230-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.