



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SCAMP5 siRNA (m): sc-153245

BACKGROUND

Secretory carrier membrane proteins (SCAMPs) are components of post Golgi membranes that are involved in endocytosis, vesicle recycling and membrane trafficking. The structural features of SCAMPs include multiple N-terminal NPF repeats and four highly conserved transmembrane regions. These NPF repeats frequently interact with EH domain proteins and aid in the budding of transport vesicles from the plasma membrane or the Golgi complex. Endocytic budding at the plasma membrane and vesicle budding at the *trans*-Golgi complex facilitates binding of SCAMP proteins to EH domain proteins. SCAMP5 (secretory carrier-associated membrane protein 5) is a 235 amino acid multi-pass membrane protein of the Golgi apparatus that exists as three alternatively spliced isoforms. Expressed in bone marrow, trachea, adrenal gland, brain, thyroid, stomach, lymph node and spinal cord, SCAMP5 plays a role in calcium-dependent exocytosis of certain cytokines, including RANTES.

REFERENCES

1. Paoluzi, S., Castagnoli, L., Lauro, I., Salcini, A.E., Coda, L., Fre', S., Confalonieri, S., Pelicci, P.G., Di Fiore, P.P. and Cesareni, G. 1998. Recognition specificity of individual EH domains of mammals and yeast. *EMBO J.* 17: 6541-6550.
2. Fernández-Chacón, R. and Südhof, T.C. 2000. Novel SCAMPs lacking NPF repeats: ubiquitous and synaptic vesicle-specific forms implicate SCAMPs in multiple membrane-trafficking functions. *J. Neurosci.* 20: 7941-7950.
3. Lin, P.J., Williams, W.P., Luu, Y., Molday, R.S., Orlowski, J. and Numata, M. 2005. Secretory carrier membrane proteins interact and regulate trafficking of the organellar (Na⁺,K⁺)/H⁺ exchanger NHE7. *J. Cell Sci.* 118: 1885-1897.
4. Noh, J.Y., Lee, H., Song, S., Kim, N.S., Im, W., Kim, M., Seo, H., Chung, C.W., Chang, J.W., Ferrante, R.J., Yoo, Y.J., Ryu, H. and Jung, Y.K. 2009. SCAMP5 links endoplasmic reticulum stress to the accumulation of expanded polyglutamine protein aggregates via endocytosis inhibition. *J. Biol. Chem.* 284: 11318-11325.
5. Han, C., Chen, T., Yang, M., Li, N., Liu, H. and Cao, X. 2009. Human SCAMP5, a novel secretory carrier membrane protein, facilitates calcium-triggered cytokine secretion by interaction with SNARE machinery. *J. Immunol.* 182: 2986-2996.
6. Castermans, D., Volders, K., Crepel, A., Backx, L., De Vos, R., Freson, K., Meulemans, S., Vermeesch, J.R., Schrandt-Stumpel, C.T., De Rijk, P., Del-Favero, J., Van Geet, C., Van De Ven, W.J., Steyaert, J.G., et al. 2010. SCAMP5, NBEA and AMISYN: three candidate genes for autism involved in secretion of large dense-core vesicles. *Hum. Mol. Genet.* 19: 1368-1378.

CHROMOSOMAL LOCATION

Genetic locus: Scamp5 (mouse) mapping to 9 B.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

SCAMP5 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SCAMP5 shRNA Plasmid (m): sc-153245-SH and SCAMP5 shRNA (m) Lentiviral Particles: sc-153245-V as alternate gene silencing products.

For independent verification of SCAMP5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153245A, sc-153245B and sc-153245C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SCAMP5 siRNA (m) is recommended for the inhibition of SCAMP5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SCAMP5 gene expression knockdown using RT-PCR Primer: SCAMP5 (m)-PR: sc-153245-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.