

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

SCUBE2 siRNA (m): sc-153276



BACKGROUND

SCUBE2 (signal peptide, CUB domain, EGF-like 2), also known as CEGP1, Cegb1 or Cegf1, is a 999 amino acid protein that is ubiquitously expressed in adult tissues and belongs to the evolutionarily conserved SCUBE protein family. Containing a CUB domain and nine EGF-like domains, SCUBE2 manifests as a secreted surface-anchored glycoprotein when overexpressed and is considered a novel component of the HH (Hedgehog) signal. The HH signal plays a pivotal role in induction of ventral neuronal and muscle cell types around the midline during vertebrate development. It is suggested that SCUBE2 expression is important in breast cancer progression and may serve as a useful prognostic marker. SCUBE2 forms homo-oligomers and heterooligomers with SCUBE1 and SCUBE3. Expressed as three isoforms produced by alternative splicing events, SCUBE2 is encoded by a gene located on human chromosome 11, which houses over 1,400 genes and comprises nearly 4% of the human genome.

REFERENCES

- 1. Grimmond, S., et al. 2001. Expression of a novel mammalian epidermal growth factor-related gene during mouse neural development. Mech. Dev. 102: 209-211.
- Yang, R.B., et al. 2002. Identification of a novel family of cell-surface proteins expressed in human vascular endothelium. J. Biol. Chem. 277: 46364-46373.
- Kawakami, A., et al. 2005. The zebrafish-secreted matrix protein you/ scube2 is implicated in long-range regulation of hedgehog signaling. Curr. Biol. 15: 480-488.
- Woods, I.G., et al. 2005. The you gene encodes an EGF-CUB protein essential for Hedgehog signaling in zebrafish. PLoS Biol. 3: e66.
- Hollway, G.E., et al. 2006. SCUBE2 mediates Hedgehog signalling in the zebrafish embryo. Dev. Biol. 294: 104-118.
- Tsai, M.T., et al. 2009. Isolation and characterization of a secreted, cellsurface glycoprotein SCUBE2 from humans. Biochem. J. 422: 119-128.
- Cheng, C.J., et al. 2009. SCUBE2 suppresses breast tumor cell proliferation and confers a favorable prognosis in invasive breast cancer. Cancer Res. 69: 3634-3641.

CHROMOSOMAL LOCATION

Genetic locus: Scube2 (mouse) mapping to 7 F1.

PRODUCT

SCUBE2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SCUBE2 shRNA Plasmid (m): sc-153276-SH and SCUBE2 shRNA (m) Lentiviral Particles: sc-153276-V as alternate gene silencing products.

For independent verification of SCUBE2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153276A, sc-153276B and sc-153276C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

 $\ensuremath{\mathsf{SCUBE2}}$ siRNA (m) is recommended for the inhibition of $\ensuremath{\mathsf{SCUBE2}}$ expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

SCUBE2 (G-4): sc-398607 is recommended as a control antibody for monitoring of SCUBE2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG K BP-FITC: sc-516140 or m-IgG K BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SCUBE2 gene expression knockdown using RT-PCR Primer: SCUBE2 (m)-PR: sc-153276-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.