



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Sec23IP siRNA (m): sc-153309

BACKGROUND

COPII-coated vesicles form on the endoplasmic reticulum by the stepwise recruitment of three cytosolic components: Sar1-GTP to initiate coat formation, Sec23/24 heterodimer to select SNARE and cargo molecules, and Sec13/31 to induce coat polymerization and membrane deformation. Sec23 is the functional human counterpart of the yeast COPII component Sec23p, which suggests that it plays a similar role in mammalian protein export from the ER. Sec23IP (Sec23 interacting protein), also known as P125 or MSTP053, is a 1,000 amino acid protein that is ubiquitously expressed and interacts with Sec23 through its N-terminus. Sec23IP is suggested to assist in the early secretory pathway as well as in the organization of exit sites of the endoplasmic reticulum. A member of the PA-PLA1 family, Sec23IP exists as two alternatively spliced isoforms containing one DDHD domain and a SAM (sterile α motif) domain.

REFERENCES

1. Ruohola, H., Kabcenell, A.K. and Ferro-Novick, S. 1988. Reconstitution of protein transport from the endoplasmic reticulum to the Golgi complex in yeast: the acceptor Golgi compartment is defective in the Sec23 mutant. *J. Cell Biol.* 107: 1465-1476.
2. Paccaud, J.P., Reith, W., Carpentier, J.L., Ravazzola, M., Amherdt, M., Schekman, R. and Orci, L. 1996. Cloning and functional characterization of mammalian homologues of the COPII component Sec23. *Mol. Biol. Cell* 7: 1535-1546.
3. Tani, K., Mizoguchi, T., Iwamatsu, A., Hatsuzawa, K. and Tagaya, M. 1999. p125 is a novel mammalian Sec23p-interacting protein with structural similarity to phospholipid-modifying proteins. *J. Biol. Chem.* 274: 20505-20512.
4. Weidler, M., Reinhard, C., Friedrich, G., Wieland, F.T. and Rösch, P. 2000. Structure of the cytoplasmic domain of p23 in solution: implications for the formation of COPI vesicles. *Biochem. Biophys. Res. Commun.* 271: 401-408.
5. Mizoguchi, T., Nakajima, K., Hatsuzawa, K., Nagahama, M., Hauri, H.P., Tagaya, M. and Tani, K. 2000. Determination of functional regions of p125, a novel mammalian Sec23p-interacting protein. *Biochem. Biophys. Res. Commun.* 279: 144-149.
6. Nakajima, K., Sonoda, H., Mizoguchi, T., Aoki, J., Arai, H., Nagahama, M., Tagaya, M. and Tani, K. 2002. A novel phospholipase A1 with sequence homology to a mammalian Sec23p-interacting protein, p125. *J. Biol. Chem.* 277: 11329-11335.
7. Bi, X., Corpina, R.A. and Goldberg, J. 2002. Structure of the Sec23/24-Sar1 pre-budding complex of the COPII vesicle coat. *Nature* 419: 271-277.
8. Duden, R. 2003. ER-to-Golgi transport: COP I and COP II function (Review). *Mol. Membr. Biol.* 20: 197-207.
9. Hughes, H. and Stephens, D.J. 2008. Assembly, organization, and function of the COPII coat. *Histochem. Cell Biol.* 129: 129-151.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Sec23ip (mouse) mapping to 7 F3.

PRODUCT

Sec23IP siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Sec23IP shRNA Plasmid (m): sc-153309-SH and Sec23IP shRNA (m) Lentiviral Particles: sc-153309-V as alternate gene silencing products.

For independent verification of Sec23IP (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153309A, sc-153309B and sc-153309C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Sec23IP siRNA (m) is recommended for the inhibition of Sec23IP expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Sec23IP gene expression knockdown using RT-PCR Primer: Sec23IP (m)-PR: sc-153309-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.