



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Selenoprotein T siRNA (m): sc-153329

BACKGROUND

Selenium is an essential trace element that is incorporated as selenocysteine into the primary structure of selenoproteins. Nutritional deficiency of selenium decreases selenoprotein concentrations and leads to pathologic conditions. Most of the known selenoproteins are members of the glutathione peroxidase or iodothyronine deiodinase families. Selenoprotein T, also known as SELT, is a 195 amino acid ubiquitously expressed protein that localizes to the endoplasmic reticulum and is a member of the SelWTH family and SELT subfamily. Selenoprotein T contains a selenocysteine (Sec) residue at its active site and is thought to be involved in redox regulation and cell anchorage. Between Cys-46 and Sec-49, Selenoprotein T may contain a selenide-sulfide bond, which is speculated to serve as the redox active pair. It is suggested that selenoprotein T regulates Ca^{2+} homeostasis and neuroendocrine secretion in response to a cAMP-stimulating trophic factor.

REFERENCES

1. Kryukov, G.V., Kryukov, V.M. and Gladyshev, V.N. 1999. New mammalian selenocysteine-containing proteins identified with an algorithm that searches for selenocysteine insertion sequence elements. *J. Biol. Chem.* 274: 33888-33897.
2. Kryukov, G.V., Castellano, S., Novoselov, S.V., Lobanov, A.V., Zehrab, O., Guigó, R. and Gladyshev, V.N. 2003. Characterization of mammalian selenoproteomes. *Science* 300: 1439-1443.
3. Dikiy, A., Novoselov, S.V., Fomenko, D.E., Sengupta, A., Carlson, B.A., Cerny, R.L., Ginalski, K., Grishin, N.V., Hatfield, D.L. and Gladyshev, V.N. 2007. SelT, SelW, SelH, and Rdx12: genomics and molecular insights into the functions of selenoproteins of a novel thioredoxin-like family. *Biochemistry* 46: 6871-6882.
4. Bukalis, K., Wolf, C., Behne, D. and Kyriakopoulos, A. 2007. Studies on the selenoproteome in the cultured cells of lung and trachea by gel electrophoretic techniques. *J. Chromatogr. A* 1155: 180-186.
5. Hoffmann, P.R. 2008. Selenium and asthma: a complex relationship. *Allergy* 63: 854-856.
6. Pappas, A.C., Zoidis, E., Surai, P.F. and Zervas, G. 2008. Selenoproteins and maternal nutrition. *Comp. Biochem. Physiol. B, Biochem. Mol. Biol.* 151: 361-372.
7. Grumolato, L., Ghzili, H., Montero-Hadjadje, M., Gasman, S., Lesage, J., Tanguy, Y., Galas, L., Ait-Ali, D., Leprince, J., Guerineau, N.C., Elkahloun, A.G., Fournier, A., Vieau, D., Vaudry, H. and Anouar, Y. 2008. Selenoprotein T is a PACAP-regulated gene involved in intracellular Ca^{2+} mobilization and neuroendocrine secretion. *FASEB J.* 22: 1756-1768.
8. Sengupta, A., Carlson, B.A., Labunsky, V.M., Gladyshev, V.N. and Hatfield, D.L. 2009. Selenoprotein T deficiency alters cell adhesion and elevates Selenoprotein W expression in murine fibroblast cells. *Biochem. Cell Biol.* 87: 953-961.
9. Shchedrina, V.A., Zhang, Y., Labunsky, V.M., Hatfield, D.L. and Gladyshev, V.N. 2010. Structure-function relations, physiological roles, and evolution of mammalian ER-resident selenoproteins. *Antioxid. Redox Signal.* 12: 839-849.

CHROMOSOMAL LOCATION

Genetic locus: Selt (mouse) mapping to 3 D.

PRODUCT

Selenoprotein T siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Selenoprotein T shRNA Plasmid (m): sc-153329-SH and Selenoprotein T shRNA (m) Lentiviral Particles: sc-153329-V as alternate gene silencing products.

For independent verification of Selenoprotein T (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153329A, sc-153329B and sc-153329C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Selenoprotein T siRNA (m) is recommended for the inhibition of Selenoprotein T expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Selenoprotein T gene expression knockdown using RT-PCR Primer: Selenoprotein T (m)-PR: sc-153329-PR (20 μ l). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.