



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

SEMA4B siRNA (m): sc-153333

BACKGROUND

Semaphorins are a family of cell surface and secreted proteins that are conserved from insects to humans. Members of this family of proteins are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular "semaphorin" domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. The transmembrane semaphorins are characterized by an additional 80 amino acid transmembrane domain and an 80-110 amino acid cytoplasmic domain. Secreted and cell-bound semaphorins chemically attract and repel the growth of neural axons, guiding the development of intricate networks of neural tissue. SEMA4B (semaphorin-4B), also known as SemC or SEMAC, is an 832 amino acid single-pass type I membrane protein that belongs to the semaphorin family and exists as two alternatively spliced isoforms. Containing one Ig-like C2-type (immunoglobulin-like) domain, a PSI domain and a single sema domain, SEMA4B is encoded by a gene located on human chromosome 15.

REFERENCES

1. Kolodkin, A.L., Matthes, D.J. and Goodman, C.S. 1993. The semaphorin genes encode a family of transmembrane and secreted growth cone guidance molecules. *Cell* 75: 1389-1399.
2. Dodd, J. and Schuchardt, A. 1995. Axon guidance: a compelling case for repelling growth cones. *Cell* 81: 471-474.
3. Matthes, D.J., Sink, H., Kolodkin, A.L. and Goodman, C.S. 1995. Semaphorin II can function as a selective inhibitor of specific synaptic arborizations. *Cell* 81: 631-639.
4. Inagaki, S., Furuyama, T. and Iwahashi, Y. 1995. Identification of a member of mouse semaphorin family. *FEBS Lett.* 370: 269-272.
5. Püschel, A.W., Adams, R.H. and Betz, H. 1995. Murine semaphorin D/collapsin is a member of a diverse gene family and creates domains inhibitory for axonal extension. *Neuron* 14: 941-948.
6. Burkhardt, C., Müller, M., Badde, A., Garner, C.C., Gundelfinger, E.D. and Püschel, A.W. 2005. Semaphorin 4B interacts with the post-synaptic density protein PSD-95/SAP90 and is recruited to synapses through a C-terminal PDZ-binding motif. *FEBS Lett.* 579: 3821-3828.
7. Nagai, H., Sugito, N., Matsubara, H., Tatematsu, Y., Hida, T., Sekido, Y., Nagino, M., Nimura, Y., Takahashi, T. and Osada, H. 2007. CLCP1 interacts with semaphorin 4B and regulates motility of lung cancer cells. *Oncogene* 26: 4025-4031.
8. Pasterkamp, R.J. and Giger, R.J. 2009. Semaphorin function in neural plasticity and disease. *Curr. Opin. Neurobiol.* 19: 263-274.

CHROMOSOMAL LOCATION

Genetic locus: Sema4b (mouse) mapping to 7 D3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

SEMA4B siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SEMA4B shRNA Plasmid (m): sc-153333-SH and SEMA4B shRNA (m) Lentiviral Particles: sc-153333-V as alternate gene silencing products.

For independent verification of SEMA4B (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153333A, sc-153333B and sc-153333C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SEMA4B siRNA (m) is recommended for the inhibition of SEMA4B expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SEMA4B gene expression knockdown using RT-PCR Primer: SEMA4B (m)-PR: sc-153333-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.