



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SERTAD2 siRNA (m): sc-153377

BACKGROUND

SERTAD2 (SERTA domain containing 2), also known as Sei-2 or TRIP-Br2, is a 314 amino acid protein that contains one SERTA domain and functions to integrate the signals provided by transcription factors at E2F-responsive promoters, thereby enhancing the effect of select transcription factors on DNA. Overexpression of SERTAD2 is thought to promote carcinogenesis, suggesting a role for SERTAD2 in tumor formation and metastasis. The gene encoding SERTAD2 maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes. Additionally, an extremely rare recessive genetic disorder, Alström syndrome, is caused by mutations in the ALMS1 gene, which maps to chromosome 2.

REFERENCES

- Nagase, T., Seki, N., Tanaka, A., Ishikawa, K. and Nomura, N. 1995. Prediction of the coding sequences of unidentified human genes. IV. The coding sequences of 40 new genes (KIAA0121-KIAA0160) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res.* 2: 167-174, 199.
- Hsu, S.I., Yang, C.M., Sim, K.G., Hentschel, D.M., O'Leary, E. and Bonventre, J.V. 2001. TRIP-Br: a novel family of PHD zinc finger- and bromodomain-interacting proteins that regulate the transcriptional activity of E2F-1/DP-1. *EMBO J.* 20: 2273-2285.
- Sim, K.G., Zang, Z., Yang, C.M., Bonventre, J.V. and Hsu, S.I. 2004. TRIP-Br links E2F to novel functions in the regulation of cyclin E expression during cell cycle progression and in the maintenance of genomic stability. *Cell Cycle* 3: 1296-1304.
- Watanabe-Fukunaga, R., Iida, S., Shimizu, Y., Nagata, S. and Fukunaga, R. 2005. SEI family of nuclear factors regulates p53-dependent transcriptional activation. *Genes Cells* 10: 851-860.
- Lai, I.L., Wang, S.Y., Yao, Y.L. and Yang, W.M. 2007. Transcriptional and subcellular regulation of the TRIP-Br family. *Gene* 388: 102-109.
- Cheong, J.K., Gunaratnam, L. and Hsu, S.I. 2008. CRM1-mediated nuclear export is required for 26 S proteasome-dependent degradation of the TRIP-Br2 proto-oncoprotein. *J. Biol. Chem.* 283: 11661-11676.
- Cheong, J.K., Gunaratnam, L., Zang, Z.J., Yang, C.M., Sun, X., Nasr, S.L., Sim, K.G., Peh, B.K., Rashid, S.B., Bonventre, J.V., Salto-Tellez, M. and Hsu, S.I. 2009. TRIP-Br2 promotes oncogenesis in nude mice and is frequently overexpressed in multiple human tumors. *J. Transl. Med.* 7: 8.

CHROMOSOMAL LOCATION

Genetic locus: Sertad2 (mouse) mapping to 11 A3.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

SERTAD2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SERTAD2 shRNA Plasmid (m): sc-153377-SH and SERTAD2 shRNA (m) Lentiviral Particles: sc-153377-V as alternate gene silencing products.

For independent verification of SERTAD2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153377A, sc-153377B and sc-153377C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SERTAD2 siRNA (m) is recommended for the inhibition of SERTAD2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SERTAD2 gene expression knockdown using RT-PCR Primer: SERTAD2 (m)-PR: sc-153377-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.