



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SETD3 siRNA (m): sc-153384

BACKGROUND

SETD3 (SET domain containing 3) is a 594 amino acid protein that belongs to the SETD3 family. Containing one SET domain and localizing to nucleus, SETD3 is a histone methyltransferase that methylates "Lys-36" of histone H3. Overexpression of the SETD3 protein in zebrafish decreases cell viability and activates caspase-3, indicating possible roles in apoptotic cell death and cell cycle regulation. In zebrafish the *setd3* protein is composed of 17 α -helices and 8 β -sheets in the secondary structures. Existing as three alternatively spliced isoforms, the SETD3 gene is conserved in chimpanzee, canine, bovine, mouse, rat, chicken, zebrafish, fruit fly, *C.elegans*, *N.crassa* and *A.thaliana*, and maps to human chromosome 14q32.2. Chromosome 14 contains about 700 genes and 106 million base pairs and makes up about 3.5% of human cellular DNA.

REFERENCES

- Heilig, R., Eckenberg, R., Petit, J.L., Fonknechten, N., Da Silva, C., Cattolico, L., Levy, M., Barbe, V., de Berardinis, V., Ureta-Vidal, A., Pelletier, E., Vico, V., Anthonard, V., Rowen, L., Madan, A., et al. 2003. The DNA sequence and analysis of human chromosome 14. *Nature* 421: 601-607.
- De Keersmaecker, K., Real, P.J., Gatta, G.D., Palomero, T., Sulis, M.L., Tosello, V., Van Vlierberghe, P., Barnes, K., Castillo, M., Sole, X., Hadler, M., Lenz, J., Aplan, P.D., Kelliher, M., Kee, B.L., Pandolfi, P.P., et al. 2010. The TLX1 oncogene drives aneuploidy in T cell transformation. *Nat. Med.* 16: 1321-1327.
- Pallister, P.D., Pallister, A.B., South, S., Toydemir, R., Johnson, J.P., Beischel, L. and Opitz, J.M. 2011. A deletion 13q34/duplication 14q32.2-14q32.33 syndrome diagnosed 50 years after neonatal presentation as infantile hypercalcemia. *Am. J. Med. Genet. A* 155A: 833-839.
- Kim, D.W., Kim, K.B., Kim, J.Y. and Seo, S.B. 2011. Characterization of a novel histone H3K36 methyltransferase SETD3 in zebrafish. *Biosci. Biotechnol. Biochem.* 75: 289-294.
- Campagna-Slater, V., Mok, M.W., Nguyen, K.T., Feher, M., Najmanovich, R. and Schapira, M. 2011. Structural chemistry of the histone methyltransferases cofactor binding site. *J. Chem. Inf. Model.* 51: 612-623.
- Petrosian, T.C. and Clarke, S.G. 2011. Uncovering the human methyltransferase. *Mol. Cell. Proteomics* 10: M110.000976.

CHROMOSOMAL LOCATION

Genetic locus: *Setd3* (mouse) mapping to 12 F1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

SETD3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SETD3 shRNA Plasmid (m): sc-153384-SH and SETD3 shRNA (m) Lentiviral Particles: sc-153384-V as alternate gene silencing products.

For independent verification of SETD3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153384A, sc-153384B and sc-153384C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SETD3 siRNA (m) is recommended for the inhibition of SETD3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SETD3 gene expression knockdown using RT-PCR Primer: SETD3 (m)-PR: sc-153384-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.