



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

SLC35B2 siRNA (m): sc-153532

BACKGROUND

Sulfation is an important post-translational modification of proteoglycans, glycolipids and glycoproteins that requires activity of 3'-phosphoadenosine 5'-phosphosulfate (PAPS), the universal sulfate donor. PAPST1 (Adenosine 3'-phospho 5'-phosphosulfate transporter 1), also known as SLC35B2 (solute carrier family 35 member B2) is a 432 amino acid type III transmembrane protein that transports PAPS from the cytosol into the Golgi apparatus. Overexpression of either PAPST1 or PAPST2, both of which are members of the nucleotide-sugar transporter family, leads to increased PAPS transport activity within the colon. In *C. elegans*, neuronal deficits correlate with reduced complexity of heparan sulfate patterns as mediated by PAPST1, suggesting that PAPST1 is required for optimum nervous system development. There are three isoforms of PAPST1 that are produced as a result of alternative splicing events.

REFERENCES

1. Kamiyama, S., Suda, T., Ueda, R., Suzuki, M., Okubo, R., Kikuchi, N., Chiba, Y., Goto, S., Toyoda, H., Saigo, K., Watanabe, M., Narimatsu, H., Jigami, Y. and Nishihara, S. 2003. Molecular cloning and identification of 3'-phosphoadenosine 5'-phosphosulfate transporter. *J. Biol. Chem.* 278: 25958-25963.
2. Kamiyama, S., Sasaki, N., Goda, E., Ui-Tei, K., Saigo, K., Narimatsu, H., Jigami, Y., Kannagi, R., Irimura, T. and Nishihara, S. 2006. Molecular cloning and characterization of a novel 3'-phosphoadenosine 5'-phosphosulfate transporter, PAPST2. *J. Biol. Chem.* 281: 10945-10953.
3. Shimazu, D., Yamamoto, N., Umino, A., Ishii, S., Sakurai, S. and Nishikawa, T. 2006. Inhibition of D-serine accumulation in the *Xenopus* oocyte by expression of the rat ortholog of human 3'-phosphoadenosine 5'-phosphosulfate transporter gene isolated from the neocortex as D-serine modulator-1. *J. Neurochem.* 96: 30-42.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610788. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Dick, G., Grøndahl, F. and Prydz, K. 2008. Overexpression of the 3'-phosphoadenosine 5'-phosphosulfate (PAPS) transporter 1 increases sulfation of chondroitin sulfate in the apical pathway of MDCK II cells. *Glycobiology* 18: 53-65.
6. Clement, A., Wiweger, M., von der Hardt, S., Rusch, M.A., Selleck, S.B., Chien, C.B. and Roehl, H.H. 2008. Regulation of zebrafish skeletogenesis by *ext2/dackel* and *papst1/pinscher*. *PLoS Genet.* 4: e1000136.
7. Bhattacharya, R., Townley, R.A., Berry, K.L. and Bülow, H.E. 2009. The PAPS transporter PST-1 is required for heparan sulfation and is essential for viability and neural development in *C. elegans*. *J. Cell Sci.* 122: 4492-4504.
8. Sasaki, N., Hirano, T., Ichimiya, T., Wakao, M., Hirano, K., Kinoshita-Toyoda, A., Toyoda, H., Suda, Y. and Nishihara, S. 2009. The 3'-phosphoadenosine 5'-phosphosulfate transporters, PAPST1 and 2, contribute to the maintenance and differentiation of mouse embryonic stem cells. *PLoS ONE* 4: e8262.

CHROMOSOMAL LOCATION

Genetic locus: *Slc35b2* (mouse) mapping to 17 B3.

PRODUCT

SLC35B2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SLC35B2 shRNA Plasmid (m): sc-153532-SH and SLC35B2 shRNA (m) Lentiviral Particles: sc-153532-V as alternate gene silencing products.

For independent verification of SLC35B2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153532A, sc-153532B and sc-153532C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SLC35B2 siRNA (m) is recommended for the inhibition of SLC35B2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SLC35B2 gene expression knockdown using RT-PCR Primer: SLC35B2 (m)-PR: sc-153532-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.