



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SLC38A7 siRNA (m): sc-153552

BACKGROUND

SLC38A7 (solute carrier family 38, member 7), also known as SNAT7, is a 462 amino acid multi-pass membrane protein that belongs to the amino acid/polyamine transporter 2 family. Members of the SLC38 family encode sodium-coupled neutral amino acid transporters (SNATs). SNATs can be further classified into system A or system N transporters based on their functional properties and patterns of substrate recognition. SLC38A7 is a system N transporter that has a substrate preference for L-glutamine. SLC38A7 also transports other amino acids with polar side chains, as well as L-histidine and L-alanine. The SLC38A7 protein is exclusively expressed in liver. Existing as two alternatively spliced isoforms, the SLC38A7 gene is conserved in chimpanzee, canine, bovine, mouse, rat, chicken and zebrafish, and maps to human chromosome 16q21. Chromosome 16 encodes over 900 genes in approximately 90 million base pairs, makes up nearly 3% of human cellular DNA and is associated with a variety of genetic disorders.

REFERENCES

1. Ben Hamida, C., Cavalier, L., Belal, S., Sanhaji, H., Nadal, N., Barhoumi, C., M'Rissa, N., Marzouki, N., Mandel, J.L., Ben Hamida, M., Koenig, M. and Hentati, F. 1997. Homozygosity mapping of giant axonal neuropathy gene to chromosome 16q24.1. *Neurogenetics* 1: 129-133.
2. Fredriksson, R., Nordström, K.J., Stephansson, O., Hägglund, M.G. and Schiöth, H.B. 2008. The solute carrier (SLC) complement of the human genome: phylogenetic classification reveals four major families. *FEBS Lett.* 582: 3811-3816.
3. Sundberg, B.E., Waag, E., Jacobsson, J.A., Stephansson, O., Rumaks, J., Svirskis, S., Alsiö, J., Roman, E., Ebendal, T., Klusa, V. and Fredriksson, R. 2008. The evolutionary history and tissue mapping of amino acid transporters belonging to solute carrier families SLC32, SLC36, and SLC38. *J. Mol. Neurosci.* 35: 179-193.
4. Newton-Cheh, C., Eijgelsheim, M., Rice, K.M., de Bakker, P.I., Yin, X., Estrada, K., Bis, J.C., Marciante, K., Rivadeneira, F., Noseworthy, P.A., Sotoodehnia, N., Smith, N.L., Rotter, J.I., Kors, J.A., Witteman, J.C., et al. 2009. Common variants at ten loci influence QT interval duration in the QTGEN Study. *Nat. Genet.* 41: 399-406.
5. Hägglund, M.G., Sreedharan, S., Nilsson, V.C., Shaik, J.H., Almkvist, I.M., Bäcklin, S., Wrangle, O. and Fredriksson, R. 2011. Identification of SLC38A7 (SNAT7) protein as a glutamine transporter expressed in neurons. *J. Biol. Chem.* 286: 20500-20511.
6. Obayashi, T. and Kinoshita, K. 2011. COXPRESdb: a database to compare gene coexpression in seven model animals. *Nucleic Acids Res.* 39: D1016-D1022.
7. Online Mendelian Inheritance in Man, OMIM™. 2011. Johns Hopkins University, Baltimore, MD. MIM Number: 614236. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Slc38a7 (mouse) mapping to 8 D1.

PRODUCT

SLC38A7 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SLC38A7 shRNA Plasmid (m): sc-153552-SH and SLC38A7 shRNA (m) Lentiviral Particles: sc-153552-V as alternate gene silencing products.

For independent verification of SLC38A7 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153552A, sc-153552B and sc-153552C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SLC38A7 siRNA (m) is recommended for the inhibition of SLC38A7 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SLC38A7 gene expression knockdown using RT-PCR Primer: SLC38A7 (m)-PR: sc-153552-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.