



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SLC6A8 siRNA (m): sc-153577

BACKGROUND

SLC6A8 (solute carrier family 6 member 8), also known as CT1 (creatine transporter 1), CRT or CRTR, is a 635 amino acid multi-pass plasma membrane protein that belongs to the sodium/neurotransporter (SNF) family. Expressed in a variety of tissues including kidney, skeletal muscle, heart, brain, prostate, testis and colon, SLC6A8 functions to transport creatine into and out of cells, specifically those found in brain and muscle tissue. Creatine, an organic acid, occurs naturally and is essential for supplying the energy needed for proper muscle and nerve function. Defects in the gene encoding SLC6A8 lead to cerebral creatine deficiency and are the cause of X-linked creatine deficiency syndrome, a rare disorder characterized by facial anomalies, seizures and mental retardation. Multiple isoforms of SLC6A8 exist due to alternative splicing events.

REFERENCES

- Salomons, G.S., van Dooren, S.J., Verhoeven, N.M., Cecil, K.M., Ball, W.S., Degrauw, T.J. and Jakobs, C. 2001. X-linked creatine-transporter gene (SLC6A8) defect: a new creatine-deficiency syndrome. *Am. J. Hum. Genet.* 68: 1497-1500.
- Rosenberg, E.H., Almeida, L.S., Kleefstra, T., deGrauw, R.S., Yntema, H.G., Bahi, N., Moraine, C., Ropers, H.H., Fryns, J.P., deGrauw, T.J., Jakobs, C. and Salomons, G.S. 2004. High prevalence of SLC6A8 deficiency in X-linked mental retardation. *Am. J. Hum. Genet.* 75: 97-105.
- Mandel, J.L. 2004. Comparative frequency of fragile-X (FMR1) and creatine transporter (SLC6A8) mutations in X-linked mental retardation. *Am. J. Hum. Genet.* 75: 730-731.
- Shojaiefard, M., Christie, D.L. and Lang, F. 2005. Stimulation of the creatine transporter SLC6A8 by the protein kinases SGK1 and SGK3. *Biochem. Biophys. Res. Commun.* 334: 742-746.
- Dodd, J.R. and Christie, D.L. 2005. Substituted cysteine accessibility of the third transmembrane domain of the creatine transporter: defining a transport pathway. *J. Biol. Chem.* 280: 32649-32654.
- Schiaffino, M.C., Bellini, C., Costabello, L., Caruso, U., Jakobs, C., Salomons, G.S. and Bonioli, E. 2005. X-linked creatine transporter deficiency: clinical description of a patient with a novel SLC6A8 gene mutation. *Neurogenetics* 6: 165-168.
- Clark, A.J., Rosenberg, E.H., Almeida, L.S., Wood, T.C., Jakobs, C., Stevenson, R.E., Schwartz, C.E. and Salomons, G.S. 2006. X-linked creatine transporter (SLC6A8) mutations in about 1% of males with mental retardation of unknown etiology. *Hum. Genet.* 119: 604-610.
- Battini, R., Chilosi, A., Mei, D., Casarano, M., Alessandri, M.G., Leuzzi, V., Ferretti, G., Tosetti, M., Bianchi, M.C. and Cioni, G. 2007. Mental retardation and verbal dyspraxia in a new patient with *de novo* creatine transporter (SLC6A8) mutation. *Am. J. Med. Genet. A* 143A: 1771-1774.
- Martínez-Muñoz, C., Rosenberg, E.H., Jakobs, C. and Salomons, G.S. 2008. Identification, characterization and cloning of SLC6A8C, a novel splice variant of the creatine transporter gene. *Gene* 418: 53-59.

CHROMOSOMAL LOCATION

Genetic locus: Slc6a8 (mouse) mapping to X A7.3.

PRODUCT

SLC6A8 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SLC6A8 shRNA Plasmid (m): sc-153577-SH and SLC6A8 shRNA (m) Lentiviral Particles: sc-153577-V as alternate gene silencing products.

For independent verification of SLC6A8 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153577A, sc-153577B and sc-153577C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SLC6A8 siRNA (m) is recommended for the inhibition of SLC6A8 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SLC6A8 gene expression knockdown using RT-PCR Primer: SLC6A8 (m)-PR: sc-153577-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.