



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SPACA3 siRNA (m): sc-153696

BACKGROUND

SPACA3 (sperm acrosome associated 3), also known as sperm lysozyme-like protein 1, lysozyme-like protein 3, lysozyme-like acrosomal sperm-specific secretory protein ALLP-17, cancer/testis antigen 54 (CT54), LYC3, SPRASA or LYZL3, is a 215 amino acid protein that participates in the fusion and adhesion of sperm and egg plasma membrane during fertilization. Identified as a novel cancer/testis antigen in hematologic malignancies, SPACA3 has the ability to elicit B-cell immune responses in patients with cancer and is considered a potential target for immunotherapy. A member of the glycosyl hydrolase 22 family which is expressed in testis, placenta and epididymis, SPACA3 exists as two alternatively spliced isoforms; SPACA3 isoform 1 is a single-pass type II membrane protein of the sperm acrosome whereas SPACA3 isoform 2 is a secreted protein.

REFERENCES

1. Welsh, I.R., Spitznagel, J.K. 1971. Distribution of lysosomal enzymes, cationic proteins, and bactericidal substances in subcellular fractions of human polymorphonuclear leukocytes. *Infect. Immun.* 4: 97-102.
2. Cramer, E.M., Breton-Gorius, J. 1987. Ultrastructural localization of lysozyme in human neutrophils by immunogold. *J. Leukoc. Biol.* 41: 242-247.
3. Ibrahim, H.R., Matsuzaki and T., Aoki, T. 2001. Genetic evidence that antibacterial activity of lysozyme is independent of its catalytic function. *FEBS Lett.* 506: 27-32.
4. Mandal, A., Klotz, K.L., Shetty, J., Jayes, F.L., Wolkowicz, M.J., Bolling, L.C., Coonrod, S.A., Black, M.B., Diekman, A.B., Haystead, T.A., Flickinger and C.J., Herr, J.C. 2003. SLLP1, a unique, intra-acrosomal, non-bacteriolytic, c lysozyme-like protein of human spermatozoa. *Biol. Reprod.* 68: 1525-1537.
5. Zhang, K., Gao, R., Zhang, H., Cai, X., Shen, C., Wu, C., Zhao and S., Yu, L. 2005. Molecular cloning and characterization of three novel lysozyme-like genes, predominantly expressed in the male reproductive system of humans, belonging to the c-type lysozyme/ α -lactalbumin family. *Biol. Reprod.* 73: 1064-1071.
6. Niyonsaba, F., Ogawa, H. 2005. Protective roles of the skin against infection: implication of naturally occurring human antimicrobial agents β -defensins, cathelicidin LL-37 and lysozyme. *J. Dermatol. Sci.* 40: 157-168.
7. Caballero, O.L., Chen, Y.T. 2009. Cancer/testis (CT) antigens: potential targets for immunotherapy. *Cancer Sci.* 100: 2014-2021.

CHROMOSOMAL LOCATION

Genetic locus: Spaca3 (mouse) mapping to 11 B5.

PRODUCT

SPACA3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SPACA3 shRNA Plasmid (m): sc-153696-SH and SPACA3 shRNA (m) Lentiviral Particles: sc-153696-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SPACA3 siRNA (m) is recommended for the inhibition of SPACA3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SPACA3 gene expression knockdown using RT-PCR Primer: SPACA3 (m)-PR: sc-153696-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.