



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

SURF-6 siRNA (m): sc-153936

BACKGROUND

SURF-6 (Surfeit locus protein-6) is a 361 amino acid protein that localizes to granular components of the nucleolus. Expressed ubiquitously with expression levels regulated during the cell cycle, SURF-6 is thought to function as a housekeeping protein that binds both RNA and DNA *in vitro* and may be involved in ribosome assembly and biosynthesis. In mice, cells lacking SURF-6 are nonviable, further implicating a role for SURF-6 in ribosome biogenesis and, possibly, proper cell cycle progression. Human SURF-6 shares structural similarity with its fish and mouse orthologs, suggesting an evolutionary conserved role between species. SURF-6 is encoded by a gene that is located on chromosome 9 in the Surfeit gene locus, which is one of the tightest gene clusters in the human genome.

REFERENCES

- Magoulas, C. and Fried, M. 1996. The SURF-6 gene of the mouse surfeit locus encodes a novel nucleolar protein. *DNA Cell Biol.* 15: 305-316.
- Magoulas, C., et al. 1998. The SURF-6 protein is a component of the nucleolar matrix and has a high binding capacity for nucleic acids *in vitro*. *Eur. J. Cell Biol.* 75: 174-183.
- Duhig, T., et al. 1998. The human Surfeit locus. *Genomics* 52: 72-78.
- Magoulas, C. and Fried, M. 2000. Isolation and genomic analysis of the human SURF-6 gene: a member of the Surfeit locus. *Gene* 243: 115-123.
- Wolff, C.M., et al. 2002. Cloning and expression of the surfeit locus member SURF-6 during embryogenesis in *Xenopus laevis*. *DNA Seq.* 13: 149-152.
- Polzikov, M., et al. 2005. Identification of an evolutionary conserved SURF-6 domain in a family of nucleolar proteins extending from human to yeast. *Biochem. Biophys. Res. Commun.* 327: 143-149.
- Gurchenkov, V.V., et al. 2005. Properties and functions of a new nucleolar protein, SURF-6, in 3T3 mouse cells. *Bioorg. Khim.* 31: 578-585.
- Romanova, L.G., et al. 2006. Implication of nucleolar protein SURF-6 in ribosome biogenesis and preimplantation mouse development. *Biol. Reprod.* 75: 690-696.
- Polzikov, M., et al. 2007. The nucleolar protein SURF-6 is essential for viability in mouse NIH/3T3 cells. *Mol. Biol. Rep.* 34: 155-160.

CHROMOSOMAL LOCATION

Genetic locus: Surf6 (mouse) mapping to 2 A3.

PRODUCT

SURF-6 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SURF-6 shRNA Plasmid (m): sc-153936-SH and SURF-6 shRNA (m) Lentiviral Particles: sc-153936-V as alternate gene silencing products.

For independent verification of SURF-6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153936A, sc-153936B and sc-153936C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SURF-6 siRNA (m) is recommended for the inhibition of SURF-6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

SURF-6 (A-5): sc-515439 is recommended as a control antibody for monitoring of SURF-6 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SURF-6 gene expression knockdown using RT-PCR Primer: SURF-6 (m)-PR: sc-153936-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.