



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SUV420H1 siRNA (m): sc-153945

BACKGROUND

SUV420H1 (suppressor of variegation 4-20 homolog 1), also known as CGI85 or KMT5B, is an 885 amino acid protein that is associated with pericentric heterochromatin in the nucleus. One of several members of the histone-lysine methyltransferase family, SUV420H1 functions as a histone methyltransferase that trimethylates the Lys-20 residue of Histone H4, thereby tagging H4 for transcriptional repression. Characteristic of most histone methyltransferases, SUV420H1 contains one SET domain through which it confers its enzymatic activity. The co-localization of SUV420H1 with pericentric heterochromatin allows the methyltransferase to play a key role in the establishment of constitutive heterochromatin, further implicating SUV420H1 as a regulator of transcriptional events. Expression of SUV420H1 is strongly down-regulated in breast cancer cells, suggesting a role for SUV420H1 in tumor suppression. Three isoforms of SUV420H1 are expressed due to alternative splicing events.

REFERENCES

- Schotta, G., et al. 2004. A silencing pathway to induce H3-K9 and H4-K20 trimethylation at constitutive heterochromatin. *Genes Dev.* 18: 1251-1262.
- Gonzalo, S., et al. 2005. Role of the RB1 family in stabilizing histone methylation at constitutive heterochromatin. *Nat. Cell Biol.* 7: 420-428.
- Pogribny, I.P., et al. 2006. Histone H3 lysine 9 and H4 lysine 20 trimethylation and the expression of Suv4-20h2 and Suv-39h1 histone methyltransferases in hepatocarcinogenesis induced by methyl deficiency in rats. *Carcinogenesis* 27: 1180-1186.
- Tryndyak, V.P., et al. 2006. Loss of DNA methylation and histone H4 lysine 20 trimethylation in human breast cancer cells is associated with aberrant expression of DNA methyltransferase 1, Suv4-20h2 histone methyltransferase and methyl-binding proteins. *Cancer Biol. Ther.* 5: 65-70.
- Benetti, R., et al. 2007. Suv4-20h deficiency results in telomere elongation and derepression of telomere recombination. *J. Cell Biol.* 178: 925-936.
- Szafrański, K., et al. 2007. Violating the splicing rules: TG dinucleotides function as alternative 3' splice sites in U2-dependent introns. *Genome Biol.* 8: R154-R154.

CHROMOSOMAL LOCATION

Genetic locus: Suv420h1 (mouse) mapping to 19 A.

PRODUCT

SUV420H1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SUV420H1 shRNA Plasmid (m): sc-153945-SH and SUV420H1 shRNA (m) Lentiviral Particles: sc-153945-V as alternate gene silencing products.

For independent verification of SUV420H1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153945A, sc-153945B and sc-153945C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SUV420H1 siRNA (m) is recommended for the inhibition of SUV420H1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SUV420H1 gene expression knockdown using RT-PCR Primer: SUV420H1 (m)-PR: sc-153945-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.