



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TCFL5 siRNA (m): sc-154137

BACKGROUND

Transcription factors containing basic helix-loop-helix (bHLH) motifs are responsible for the regulation of neurogenesis, cardiogenesis, myogenesis, differentiation and cell proliferation. TCFL5 [transcription factor-like 5 (basic helix-loop-helix)], also known as cha transcription factor, HPV-16 E2-binding protein 1, CHA, Figlb, E2BP-1 or MGC46135, is a 500 amino acid protein belonging to the basic helix-loop-helix (bHLH) family. TCFL5 localizes to the nucleus and may function as a transcription factor during early spermatogenesis. TCFL5 is known to form a complex with upstream stimulatory factor (USF)-1, thereby negatively regulating USF-dependent transcription. Six isoforms of TCFL5 exist due to alternative splicing; isoform 2 is pancreas specific and isoform 3 is testis specific showing expression in primary spermatocytes at the pachytene stage. Antibodies against TCFL5 have been identified in patients with Chagas' disease, which is caused by *Trypanosoma cruzi* infection.

REFERENCES

- Gonenne, A. and Lebowitz, J. 1975. Disc electrophoretic studies of hemoglobin dissociation by p-hydroxymercury benzoate. *Anal. Biochem.* 64: 425-437.
- Maruyama, O., Nishimori, H., Katagiri, T., Miki, Y., Ueno, A. and Nakamura, Y. 1998. Cloning of TCFL5 encoding a novel human basic helix-loop-helix motif protein that is specifically expressed in primary spermatocytes at the pachytene stage. *Cytogenet. Cell Genet.* 82: 41-45.
- Online Mendelian Inheritance in Man, OMIM[™]. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 604745. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Gironès, N., Rodríguez, C.I., Carrasco-Marín, E., Hernáez, R.F., de Rego, J.L. and Fresno, M. 2001. Dominant T- and B-cell epitopes in an autoantigen linked to Chagas' disease. *J. Clin. Invest.* 107: 985-993.
- Rodríguez, C.I., Gironès, N. and Fresno, M. 2003. Cha, a basic helix-loop-helix transcription factor involved in the regulation of upstream stimulatory factor activity. *J. Biol. Chem.* 278: 43135-43145.
- Gironès, N., Cuervo, H. and Fresno, M. 2005. *Trypanosoma cruzi*-induced molecular mimicry and Chagas' disease. *Curr. Top. Microbiol. Immunol.* 296: 89-123.

CHROMOSOMAL LOCATION

Genetic locus: *Tcf15* (mouse) mapping to 2 H4.

PRODUCT

TCFL5 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCFL5 shRNA Plasmid (m): sc-154137-SH and TCFL5 shRNA (m) Lentiviral Particles: sc-154137-V as alternate gene silencing products.

For independent verification of TCFL5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154137A, sc-154137B and sc-154137C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TCFL5 siRNA (m) is recommended for the inhibition of TCFL5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCFL5 gene expression knockdown using RT-PCR Primer: TCFL5 (m)-PR: sc-154137-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.