



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TCP-11 siRNA (m): sc-154143

BACKGROUND

TCP-11 (t-complex protein 11), also known as D6S230E, is a 503 amino acid single-pass membrane protein expressed only in fertile adult testis and is a member of the TCP-11 family. Localized to the surface of mature epididymal spermatozoa, TCP-11 may be a receptor for the fertilization promoting peptide (FPP), a peptide produced by the prostate gland and then secreted into seminal plasma. The adenylate cyclase/cyclic AMP pathway is considered to be the signal transduction pathway that is activated by the association between FPP and TCP-11. TCP-11 is suggested to play a critical role in the regulation of sperm function and fertility. The gene encoding TCP-11 is located on human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Three isoforms of TCP-11 are produced by alternative splicing events.

REFERENCES

- Ragoussis, J., Senger, G., Mockridge, I., Sanseau, P., Ruddy, S., Dudley, K., Sheer, D. and Trowsdale, J. 1992. A testis-expressed Zn finger gene (ZNF76) in human 6p21.3 centromeric to the MHC is closely linked to the human homolog of the t-complex gene tcp-11. *Genomics* 14: 673-679.
- Hosseini, R., Ruddy, S., Bains, S., Hynes, G., Marsh, P., Pizze, J. and Dudley, K. 1994. The mouse t-complex gene, Tcp-11, is under translational control. *Mech. Dev.* 47: 73-80.
- Fraser, L.R., Hosseini, R., Hanyalogou, A., Talmor, A. and Dudley, R.K. 1997. TCP-11, the product of a mouse t-complex gene, plays a role in stimulation of capacitation and inhibition of the spontaneous acrosome reaction. *Mol. Reprod. Dev.* 48: 375-382.
- Fraser, L.R. 1998. The modulation of sperm function by fertilization promoting peptide. *Hum. Reprod.* 13: 1-10.
- Adeoya-Osiguwa, S.A., Dudley, R.K., Hosseini, R. and Fraser, L.R. 1998. FPP modulates mammalian sperm function via TCP-11 and the adenyl cyclase/cAMP pathway. *Mol. Reprod. Dev.* 51: 468-476.
- Fraser, L.R. and Adeoya-Osiguwa, S. 1999. Modulation of adenyl cyclase by FPP and adenosine involves stimulatory and inhibitory adenosine receptors and g proteins. *Mol. Reprod. Dev.* 53: 459-471.
- Ma, Y., Zhang, S., Xia, Q., Zhang, G., Huang, X., Huang, M., Xiao, C., Pan, A., Sun, Y., Lebo, R. and Milunsky, A. 2002. Molecular characterization of the TCP11 gene which is the human homologue of the mouse gene encoding the receptor of fertilization promoting peptide. *Mol. Hum. Reprod.* 8: 24-31.
- Safronova, L.D., Kudriavtsev, I.V. and Kudriavtsev, P.I. 2002. Sterility of males determined by functional features of the mouse spermatozoa bearing t-complex. *Ontogenez* 33: 165-169.
- Ma, Y.X., Zhang, S.Z., Wu, Q.Q., Sun, Y., Qiu, W.M. and Xu, W.M. 2003. Cloning, expression, and alternative splicing of the novel isoform of hTCP11 gene. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao* 25: 122-128.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Tcp11 (mouse) mapping to 17 A3.3.

PRODUCT

TCP-11 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCP-11 shRNA Plasmid (m): sc-154143-SH and TCP-11 shRNA (m) Lentiviral Particles: sc-154143-V as alternate gene silencing products.

For independent verification of TCP-11 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154143A and sc-154143B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TCP-11 siRNA (m) is recommended for the inhibition of TCP-11 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCP-11 gene expression knockdown using RT-PCR Primer: TCP-11 (m)-PR: sc-154143-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.