



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TDRD7 siRNA (m): sc-154168

BACKGROUND

TDRD7 (tudor domain containing 7), also known as TRAP, CATC4 or PCTAIRE2BP, is a 1,098 amino acid cytoplasmic protein that localizes to RNA granules. TDRD7 is present in the chromatoid body (CB) of spermatids and is detected in the intermitochondrial cementin of pachytene spermatocytes. TDRD7 contains three lotus/OST-HTH domains and two tudor domains, and forms a mRNP complex with TDRD1, TDRD6 and VASA. TDRD7 is a component of cytoplasmic RNA granules involved in post-transcriptional regulation of specific genes. Required for lens transparency during lens development, TDRD7 is also required during spermatogenesis. Mutations in the gene encoding TDRD7 leads to cataract congenital autosomal recessive type 4 (CATC4), which is characterized by an opacification of the crystalline lens of the eye. TDRD7 exists as three alternatively spliced isoforms and is encoded by a gene located on human chromosome 9q22.33.

REFERENCES

- Hirose, T., Kawabuchi, M., Tamaru, T., Okumura, N., Nagai, K. and Okada, M. 2000. Identification of tudor repeat associator with PCTAIRE 2 (Trap). A novel protein that interacts with the N-terminal domain of PCTAIRE 2 in rat brain. *Eur. J. Biochem.* 267: 2113-2121.
- Yamochi, T., Nishimoto, I., Okuda, T. and Matsuoka, M. 2001. ik3-1/Cables is associated with Trap and Pctaire2. *Biochem. Biophys. Res. Commun.* 286: 1045-1050.
- Hosokawa, M., Shoji, M., Kitamura, K., Tanaka, T., Noce, T., Chuma, S. and Nakatsuji, N. 2007. Tudor-related proteins TDRD1/MTR-1, TDRD6 and TDRD7/TRAP: domain composition, intracellular localization, and function in male germ cells in mice. *Dev. Biol.* 301: 38-52.
- Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611258. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Skorokhod, O., Nemazanyy, I., Breus, O., Filonenko, V. and Panasyuk, G. 2008. Generation and characterization of monoclonal antibodies to TDRD7 protein. *Hybridoma* 27: 211-216.
- Callebaut, I. and Mornon, J.P. 2010. LOTUS, a new domain associated with small RNA pathways in the germline. *Bioinformatics* 26: 1140-1144.
- Anantharaman, V., Zhang, D. and Aravind, L. 2010. OST-HTH: a novel predicted RNA-binding domain. *Biol. Direct.* 5: 13.
- Tanaka, T., Hosokawa, M., Vagin, V.V., Reuter, M., Hayashi, E., Mochizuki, A.L., Kitamura, K., Yamanaka, H., Kondoh, G., Okawa, K., Kuramochi-Miyagawa, S., Nakano, T., Sachidanandam, R., et al. 2011. Tudor domain containing 7 (Tdrd7) is essential for dynamic ribonucleoprotein (RNP) remodeling of chromatoid bodies during spermatogenesis. *Proc. Natl. Acad. Sci. USA* 108: 10579-10584.
- Lachke, S.A., Alkuraya, F.S., Kneeland, S.C., Ohn, T., Aboukhalil, A., Howell, G.R., Saadi, I., Cavallalesco, R., Yue, Y., Tsai, A.C., Nair, K.S., Cosma, M.I., Smith, R.S., Hodges, E., Alfaradhi, S.M., Al-Hajeri, A., et al. 2011. Mutations in the RNA granule component TDRD7 cause cataract and glaucoma. *Science* 331: 1571-1576.

CHROMOSOMAL LOCATION

Genetic locus: Tdrd7 (mouse) mapping to 4 B1.

PRODUCT

TDRD7 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TDRD7 shRNA Plasmid (m): sc-154168-SH and TDRD7 shRNA (m) Lentiviral Particles: sc-154168-V as alternate gene silencing products.

For independent verification of TDRD7 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154168A, sc-154168B and sc-154168C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TDRD7 siRNA (m) is recommended for the inhibition of TDRD7 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TDRD7 gene expression knockdown using RT-PCR Primer: TDRD7 (m)-PR: sc-154168-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.