



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TEF siRNA (m): sc-154177

BACKGROUND

TEF (thyrotrophic embryonic factor), also known as KIAA1655, is a 303 amino acid nuclear transcription factor that belongs to the bZIP (basic region/leucine zipper) family and PAR (proline and acidic amino acid-rich) subfamily. TEF binds DNA as either a homodimer or heterodimer, and is known to transactivate the TSH β promoter. While broadly expressed in adults, TEF is only found in developing embryonic anterior pituitary gland. TEF accumulates according to a robust circadian rhythm and has also been found to inhibit cell growth by down-regulating β chain expression of cytokine receptors. The functional domains of TEF are highly homologous with other members of the PAR-bZIP subfamily, including albumin D box-binding protein (DABP), human hepatic leukemia factor (HLF) and chicken vitellogenin gene-binding protein (VBP). The gene encoding TEF maps to human chromosome 22q13.2.

REFERENCES

1. Drolet, D.W., Scully, K.M., Simmons, D.M., Wegner, M., Chu, K.T., Swanson, L.W. and Rosenfeld, M.G. 1991. TEF, a transcription factor expressed specifically in the anterior pituitary during embryogenesis, defines a new class of leucine zipper proteins. *Genes Dev.* 5: 1739-1753.
2. Lossie, A.C., Gordon, D.F. and Camper, S.A. 1993. Localization of thyrotropin-releasing hormone receptor and thyrotrophic embryonic factor on mouse chromosome 15. *Mamm. Genome* 4: 621-623.
3. Khatib, Z.A., Inaba, T., Valentine, M. and Look, A.T. 1994. Chromosomal localization and cDNA cloning of the human DBP and TEF genes. *Genomics* 23: 344-351.
4. Krueger, D.A., Warner, E.A. and Dowd, D.R. 2000. Involvement of thyrotrophic embryonic factor in calcium-mediated regulation of gene expression. *J. Biol. Chem.* 275: 14524-14531.
5. Newman, J.R. and Keating, A.E. 2003. Comprehensive identification of human bZIP interactions with coiled-coil arrays. *Science* 300: 2097-2101.
6. Gachon, F., Fonjallaz, P., Damiola, F., Gos, P., Kodama, T., Zakany, J., Duboule, D., Petit, B., Tafti, M. and Schibler, U. 2004. The loss of circadian PAR bZip transcription factors results in epilepsy. *Genes Dev.* 18: 1397-1412.
7. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 188595. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Inukai, T., Inaba, T., Dang, J., Kuribara, R., Ozawa, K., Miyajima, A., Wu, W., Look, A.T., Arinobu, Y., Iwasaki, H., Akashi, K., Kagami, K., Goi, K., Sugita, K. and Nakazawa, S. 2005. TEF, an antiapoptotic bZIP transcription factor related to the oncogenic E2A-HLF chimera, inhibits cell growth by down-regulating expression of the common β chain of cytokine receptors. *Blood* 105: 4437-4444.

CHROMOSOMAL LOCATION

Genetic locus: Tef (mouse) mapping to 15 E1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

TEF siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TEF shRNA Plasmid (m): sc-154177-SH and TEF shRNA (m) Lentiviral Particles: sc-154177-V as alternate gene silencing products.

For independent verification of TEF (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154177A, sc-154177B and sc-154177C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TEF siRNA (m) is recommended for the inhibition of TEF expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TEF gene expression knockdown using RT-PCR Primer: TEF (m)-PR: sc-154177-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.