



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TELO2 siRNA (m): sc-154186

BACKGROUND

TELO2 (telomere maintenance 2), also known as CLK2 or TEL2, is an 837 amino acid protein that is expressed in the cytoplasm and the nucleus. Belonging to the TEL2 family, TELO2 may be involved in telomere length and growth regulation. Cells overexpressing TELO2 are hypersensitive to hydroxyurea (HU) and undergo apoptotic death in response to HU treatment. TELO2 functions as an S-phase checkpoint protein in the cell cycle and is required for survival of replication fork arrest. Considered a highly conserved regulator of PIKKs (phosphatidylinositol 3-kinase-related protein kinases), which include SMG1, TELO2 may be essential in embryonic development and may also play a role in DNA repair. The gene encoding TELO2 is located on human chromosome 16, which encodes over 900 genes and comprises nearly 3% of the human genome.

REFERENCES

1. Takai, H., et al. 2007. TEL2 regulates the stability of PI3K-related protein kinases. *Cell* 131: 1248-1259.
2. Kanoh, J., et al. 2007. TEL2: a common partner of PIK-related kinases and a link between DNA checkpoint and nutritional response? *Genes Cells* 12: 1301-1304.
3. Shikata, M., et al. 2007. TEL2 is required for activation of the Mrc1-mediated replication checkpoint. *J. Biol. Chem.* 282: 5346-5355.
4. Seidel, J.J., et al. 2008. A novel Tel1/ATM N-terminal motif, TAN, is essential for telomere length maintenance and a DNA damage response. *Mol. Cell. Biol.* 28: 5736-5746.
5. Onitake, Y., et al. 2009. Telomere biology in neuroblastoma: telomere binding proteins and alternative strengthening of telomeres. *J. Pediatr. Surg.* 44: 2258-2266.
6. Ueno, M. 2010. Roles of DNA repair proteins in telomere maintenance. *Biosci. Biotechnol. Biochem.* 74: 1-6.
7. Pennarun, G., et al. 2010. ATR contributes to telomere maintenance in human cells. *Nucleic Acids Res.* 38: 2955-2963.

CHROMOSOMAL LOCATION

Genetic locus: Telo2 (mouse) mapping to 17 A3.3.

PRODUCT

TELO2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TELO2 shRNA Plasmid (m): sc-154186-SH and TELO2 shRNA (m) Lentiviral Particles: sc-154186-V as alternate gene silencing products.

For independent verification of TELO2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154186A, sc-154186B and sc-154186C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TELO2 siRNA (m) is recommended for the inhibition of TELO2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TELO2 gene expression knockdown using RT-PCR Primer: TELO2 (m)-PR: sc-154186-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Yorichika, N., et al. 2019. The effects of TEL2 on cardiomyocyte survival. *Life Sci.* 16: 116665.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.