



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Tenascin-N siRNA (m): sc-154188

BACKGROUND

The Tenascin family of extracellular matrix proteins includes Tenascin (also designated cytotactin or Tenascin-C), Tenascin-R (also designated restrictin or janusin), Tenascin-X, and Tenascin-N (also designated TNN, TN-W or Tenascin-W). Tenascin proteins function as substrate-adhesion molecules (SAMs) and are involved in regulating numerous developmental processes, such as morphogenetic cell migration and organogenesis. The Tenascin family proteins arise from various splicing events in the region coding for fibronectin (FN) III repeats. Tenascin-C and Tenascin-X are expressed in several tissues during embryogenesis, and in adult tissues undergoing active remodeling such as healing wounds and tumors. Tenascin-R (TN-R) is expressed on the surface of neurons and glial cells and Tenascin-N, although evident during development, is predominantly expressed by neurons in the adult central nervous system. Tenascin-N may play a role in neurite outgrowth and migration functioning as a repulsive molecule in the hippocampus.

REFERENCES

- Jung, M., Pesheva, P., Schachner, M. and Trotter, J. 1993. Astrocytes and neurons regulate the expression of the neural recognition molecule janusin by cultured oligodendrocytes. *Glia* 9: 163-175.
- Schachner, M., Taylor, J., Bartsch, U. and Pesheva, P. 1994. The perplexing multifunctionality of janusin, a tenascin-related molecule. *Perspect. Dev. Neurobiol.* 2: 33-41.
- Chiquet-Ehrismann, R. 1995. Tenascins, a growing family of extracellular matrix proteins. *Experientia* 51: 853-862.
- Faissner, A. 1997. The tenascin gene family in axon growth and guidance. *Cell Tissue Res.* 290: 331-341.
- Efleteriou, F., Exposito, J.Y., Garrone, R. and Lethias, C. 1997. Characterization of the bovine Tenascin-X. *J. Biol. Chem.* 272: 22866-22874.
- Srinivasan, J., Schachner, M. and Catterall, W.A. 1998. Interaction of voltage-gated sodium channels with the extracellular matrix molecules Tenascin-C and Tenascin-R. *Proc. Natl. Acad. Sci. USA* 95: 15753-15757.
- Neidhardt, J., Fehr, S., Kutsche, M., Löhler, J. and Schachner, M. 2003. Tenascin-N: characterization of a novel member of the tenascin family that mediates neurite repulsion from hippocampal explants. *Mol. Cell. Neurosci.* 23: 193-209.
- Tucker, R.P., Drabikowski, K., Hess, J.F., Ferralli, J., Chiquet-Ehrismann, R. and Adams, J.C. 2006. Phylogenetic analysis of the tenascin gene family: evidence of origin early in the chordate lineage. *BMC Evol. Biol.* 6: 60.
- Nishida, E., Sasaki, T., Ishikawa, S.K., Kosaka, K., Aino, M., Noguchi, T., Teranaka, T., Shimizu, N. and Saito, M. 2007. Transcriptome database KK-periome for periodontal ligament development: expression profiles of the extracellular matrix genes. *Gene* 404: 70-79.

CHROMOSOMAL LOCATION

Genetic locus: Tnn (mouse) mapping to 1 H2.1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Tenascin-N siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Tenascin-N shRNA Plasmid (m): sc-154188-SH and Tenascin-N shRNA (m) Lentiviral Particles: sc-154188-V as alternate gene silencing products.

For independent verification of Tenascin-N (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154188A, sc-154188B and sc-154188C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Tenascin-N siRNA (m) is recommended for the inhibition of Tenascin-N expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Tenascin-N gene expression knockdown using RT-PCR Primer: Tenascin-N (m)-PR: sc-154188-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.