



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Titin siRNA (m): sc-154286

BACKGROUND

Titin, also known as connectin, is a large protein involved in the temporal and spatial control of the assembly of the highly ordered sarcomeres (contractile units) of striated muscle. In addition to sarcomere assembly Titin also functions to maintain the structural integrity of the contracting myofibrils within the muscle as well as organizing the machinery for condensation of chromosomes in dividing cells. Titin is a giant protein composed of 27,000 amino acids and contains an autoregulated serine kinase catalytic domain as well as a calcium/calmodulin binding region that are involved in its activation. Activated Titin phosphorylates the muscle protein Telethonin, a sarcomeric protein abundant in heart and skeletal muscle, implicating Titin activity in the reorganization of the cytoskeleton during myofibrillogenesis.

REFERENCES

1. Trinick, J. 1996. Titin as a scaffold and spring. *Cytoskeleton Curr. Biol.* 6: 258-260.
2. Valle, G., et al. 1997. Telethonin, a novel sarcomeric protein of heart and skeletal muscle. *FEBS Lett.* 415: 163-168.
4. Means, A.R. 1998. The clash in Titin. *Nature* 395: 846-847.
3. Mayans, O., et al. 1998. Structural basis for activation of the Titin kinase domain during myofibrillogenesis. *Nature* 395: 863-869. Published erratum appears in *Nature*. 1999. 397: 719.
5. Gregorio, C.C., et al. 1999. Muscle assembly: a titanic achievement? *Curr. Opin. Cell. Biol.* 11: 18-25.
6. Trinick, J. et al. 1999. Titin: a molecular control freak. *Trends Cell. Biol.* 9: 377-380.
7. Niederlander, N., et al. 2004. Regulation of the Actin-Myosin interaction by Titin. *Eur. J. Biochem.* 271: 4572-4581.
8. Fukuda, N., et al. 2005. Phosphorylation of Titin modulates passive stiffness of cardiac muscle in a Titin isoform-dependent manner. *J. Gen. Physiol.* 125: 257-271.
9. Harris, B.N., et al. 2005. Calcium transients regulate Titin organization during myofibrillogenesis. *Cell Motil. Cytoskeleton* 60: 129-139.

CHROMOSOMAL LOCATION

Genetic locus: Ttn (mouse) mapping to 2 C3.

PRODUCT

Titin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Titin shRNA Plasmid (m): sc-154286-SH and Titin shRNA (m) Lentiviral Particles: sc-154286-V as alternate gene silencing products.

For independent verification of Titin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154286A, sc-154286B and sc-154286C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Titin siRNA (m) is recommended for the inhibition of Titin expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Titin (E-2): sc-271946 is recommended as a control antibody for monitoring of Titin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Titin gene expression knockdown using RT-PCR Primer: Titin (m)-PR: sc-154286-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.