



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TMC8 siRNA (m): sc-154320

## BACKGROUND

Transmembrane channel-like protein 8 (TMC8), also known as Epidermodysplasia verruciformis protein 2 (EVER2), is a 726 amino acid member of the TMC family of proteins. Localized to the endoplasmic reticulum membrane, TMC8 is thought to form a transmembrane channel-like protein with eight predicted transmembrane domains and three leucine zipper motifs. Mutations in the genes encoding TMC8 and TMC6, another member of the TMC family, have been shown to cause epidermodysplasia verruciformis (EV), an autosomal recessive dermatosis characterized by abnormal susceptibility to human papillomaviruses (HPVs) and a high rate of progression to squamous cell carcinoma on sun-exposed skin. Infection by HPVs lead to persistent wart-like or macular lesions. TMC8 is expressed in placenta, prostate and testis, and three named isoforms exist as a result of alternative splicing events.

## REFERENCES

1. Ramoz, N., Rueda, L.A., Bouadjar, B., Montoya, L.S., Orth, G. and Favre, M. 2002. Mutations in two adjacent novel genes are associated with epidermodysplasia verruciformis. *Nat. Genet.* 32: 579-581.
2. Keresztes, G., Mutai, H. and Heller, S. 2003. TMC and EVER genes belong to a larger novel family, the TMC gene family encoding transmembrane proteins. *BMC Genomics* 4: 24.
3. Kurima, K., Yang, Y., Sorber, K. and Griffith, A.J. 2003. Characterization of the transmembrane channel-like (TMC) gene family: functional clues from hearing loss and epidermodysplasia verruciformis. *Genomics* 82: 300-308.
4. Azzimonti, B., Mondini, M., De Andrea, M., Gioia, D., Dianzani, U., Mesturini, R., Leigh, G., Tiberio, R., Landolfo, S. and Gariglio, M. 2005. CD8<sup>+</sup> T-cell lymphocytopenia and lack of EVER mutations in a patient with clinically and virologically typical epidermodysplasia verruciformis. *Arch Dermatol.* 141: 1323-1325.
5. Sun, X.K., Chen, J.F. and Xu, A.E. 2005. A homozygous nonsense mutation in the EVER2 gene leads to epidermodysplasia verruciformis. *Clin. Exp. Dermatol.* 30: 573-574.
6. Orth, G. 2006. Genetics of epidermodysplasia verruciformis: insights into host defense against papillomaviruses. *Semin. Immunol.* 18: 362-374.
7. Orth, G. 2008. Host defenses against human papillomaviruses: lessons from epidermodysplasia verruciformis. *Curr. Top. Microbiol. Immunol.* 321: 59-83.
8. Patel, A.S., Karagas, M.R., Pawlita, M., Waterboer, T. and Nelson, H.H. 2008. Cutaneous human papillomavirus infection, the EVER2 gene and incidence of squamous cell carcinoma: a case-control study. *Int. J. Cancer* 122: 2377-2379.
9. Hohenstein, E., Rady, P.L., Hergersberg, M., Huber, A.R., Tyring, S.K., Bregenzner, T., Streit, M. and Itin, P. 2009. Epidermodysplasia verruciformis in a HIV-positive patient homozygous for the c917A→T polymorphism in the TMC8/EVER2 gene. *Dermatology* 218: 114-118.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: *Tmc8* (mouse) mapping to 11 E2.

## PRODUCT

TMC8 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TMC8 shRNA Plasmid (m): sc-154320-SH and TMC8 shRNA (m) Lentiviral Particles: sc-154320-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

TMC8 siRNA (m) is recommended for the inhibition of TMC8 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TMC8 gene expression knockdown using RT-PCR Primer: TMC8 (m)-PR: sc-154320-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.