



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TMEM154 siRNA (m): sc-154386

BACKGROUND

TMEM154 is a 128 amino acid protein encoded by a gene mapping to human chromosome 4. Representing approximately 6% of the human genome, chromosome 4 contains nearly 900 genes. Notably, the Huntingtin gene, which is found to encode an expanded glutamine tract in cases of Huntington's disease, is on chromosome 4. FGFR-3 is also encoded on chromosome 4 and has been associated with thanatophoric dwarfism, achondroplasia, Muenke syndrome and bladder cancer. Chromosome 4 is also tied to Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease. Chromosome 4 reportedly contains the largest gene deserts (regions of the genome with no protein encoding genes) and has one of the two lowest recombination frequencies of the human chromosomes.

REFERENCES

- Hillier, L.W., et al. 2005. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. *Nature* 434: 724-731.
- Cowan, C.M. and Raymond, L.A. 2006. Selective neuronal degeneration in Huntington's disease. *Curr. Top. Dev. Biol.* 75:25-71.
- Chandler, R.J., et al. 2007. Metabolic phenotype of methylmalonic acidemia in mice and humans: the role of skeletal muscle. *BMC Med. Genet.* 8: 64
- Cunningham, M.L., et al. 2007. Syndromic craniosynostosis: from history to hydrogen bonds. *Orthod. Craniofac. Res.* 10: 67-81.
- de Frutos, C.A., et al. 2007. Snail1 is a transcriptional effector of FGFR-3 Signaling during chondrogenesis and achondroplasias. *Dev. Cell* 13: 872-883.
- Doherty, E.S., et al. 2007. Muenke syndrome (FGFR-3-related craniosynostosis): Expansion of the phenotype and review of the literature. *Am. J. Med. Genet. A* 143: 3204-3215.
- Ruiz-Perez, V.L., et al. 2007. EVC is a positive mediator of Ihh-regulated bone growth that localises at the base of chondrocyte cilia. *Development* 134: 2903-2912.

CHROMOSOMAL LOCATION

Genetic locus: Tmem154 (mouse) mapping to 3 F1.

PRODUCT

TMEM154 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TMEM154 shRNA Plasmid (m): sc-154386-SH and TMEM154 shRNA (m) Lentiviral Particles: sc-154386-V as alternate gene silencing products.

For independent verification of TMEM154 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154386A, sc-154386B and sc-154386C.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TMEM154 siRNA (m) is recommended for the inhibition of TMEM154 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TMEM154 gene expression knockdown using RT-PCR Primer: TMEM154 (m)-PR: sc-154386-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.