



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Tom34 siRNA (m): sc-154551

## BACKGROUND

The mitochondrial preprotein translocases of the outer membrane (Tom) form a multisubunit complex that facilitates the import of nuclear-encoded precursor proteins across the mitochondrial outer membrane. The Tom machinery consists of import receptors for the initial binding of cytosolically synthesized preproteins, and a general import pore (GIP) for the membrane translocation of various preproteins into the mitochondrion. Tom34 (translocase of outer mitochondrial membrane 34), also known as TOMM34, URCC3 or HTOM34P, is a ubiquitously expressed 309 amino acid protein that contains six TPR repeats. Localized to the cytoplasmic side of the mitochondrion, Tom34 functions as a chaperone-like protein that binds the mature portion of unfolded proteins and guides their import into mitochondria. In addition, Tom34 is thought to have weak ATPase activity and may interact with other organelles within the cell. Expression of Tom34 is upregulated in colon cancer cells and is thought to be involved in tumor progression. Tom34 may, therefore, be a novel target for therapeutic anti-cancer drugs.

## REFERENCES

- Nuttall, S.D., Hanson, B.J., Mori, M. and Hoogenraad, N.J. 1997. HTom34: a novel translocase for the import of proteins into human mitochondria. *DNA Cell Biol.* 16: 1067-1074.
- Chewawiwat, N., Yano, M., Terada, K., Hoogenraad, N.J. and Mori, M. 1999. Characterization of the novel mitochondrial protein import component, Tom34, in mammalian cells. *J. Biochem.* 125: 721-727.
- Mukhopadhyay, A., Avramova, L.V. and Weiner, H. 2002. Tom34 unlike Tom20 does not interact with the leader sequences of mitochondrial precursor proteins. *Arch. Biochem. Biophys.* 400: 97-104.
- Yang, C.S. and Weiner, H. 2002. Yeast two-hybrid screening identifies binding partners of human Tom34 that have ATPase activity and form a complex with Tom34 in the cytosol. *Arch. Biochem. Biophys.* 400: 105-110.
- Blesa, J.R., Prieto-Ruiz, J.A. and Hernández-Yago, J. 2004. Conformation-sensitive gel electrophoresis as an ideal high-throughput strategy for accurate detection of sequence variations in DNA: screening hTomm and hTimm genes. *J. Biomol. Screen.* 9: 621-624.
- Olsen, J.V., Blagoev, B., Gnad, F., Macek, B., Kumar, C., Mortensen, P. and Mann, M. 2006. Global, *in vivo*, and site-specific phosphorylation dynamics in signaling networks. *Cell* 127: 635-648.
- Shimokawa, T., Matsushima, S., Tsunoda, T., Tahara, H., Nakamura, Y. and Furukawa, Y. 2006. Identification of TOMM34, which shows elevated expression in the majority of human colon cancers, as a novel drug target. *Int. J. Oncol.* 29: 381-386.
- Blesa, J.R., Prieto-Ruiz, J.A., Abraham, B.A., Harrison, B.L., Hegde, A.A. and Hernández-Yago, J. 2008. NRF-1 is the major transcription factor regulating the expression of the human TOMM34 gene. *Biochem. Cell Biol.* 86: 46-56.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Tomm34 (mouse) mapping to 2 H3.

## PRODUCT

Tom34 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Tom34 shRNA Plasmid (m): sc-154551-SH and Tom34 shRNA (m) Lentiviral Particles: sc-154551-V as alternate gene silencing products.

For independent verification of Tom34 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154551A, sc-154551B and sc-154551C.

## APPLICATIONS

Tom34 siRNA (m) is recommended for the inhibition of Tom34 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Tom34 gene expression knockdown using RT-PCR Primer: Tom34 (m)-PR: sc-154551-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.