



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TReP-132 siRNA (m): sc-154633

BACKGROUND

TReP-132 (transcriptional-regulating factor 1, Breast cancer anti-estrogen resistance 2) is a 1,200 amino acid nuclear protein that contains three C₂H₂-type zinc fingers, one ELM2 domain, and one SANT domain. TReP-132 is believed to activate transcription of CYP11A1. TReP-132 interaction with CREBBP and EP300 results in a synergistic transcriptional activation of CYP11A1. TReP-132 was initially identified as a regulator of steroidogenesis but is also believed to be a cell growth suppressor. TReP-132 acts by inducing the gene expression of the G₁ cyclin-dependent kinase inhibitors p21^{WAF1/Cip1} (p21) and p27^{Kip1} (p27). This interaction is believed to be achieved with progesterone-bound PR (progesterone receptor) to trans activate the p21 and p27 gene promoters at proximal Sp1-binding sites. Highest expression of TReP-132 is believed to be in thymus, testis and adrenal cortex, but may also be expressed in the adrenal medulla, thyroid, and stomach. TReP-132 is highly expressed in steroidogenic JEG-3 and MCF-7 cells with low expression in non-steroidogenic Hep G2 and HK293 cells.

REFERENCES

- Gizard, F., Lavallee, B., DeWitte, F. and Hum, D.W. 2001. A novel zinc finger protein TReP-132 interacts with CBP/p300 to regulate human CYP11A1 gene expression. *J. Biol. Chem.* 276: 33881-33892.
- Gizard, F., El-Alfy, M., Duguay, Y., Lavallee, B., DeWitte, F., Staels, B., Beatty, B.G. and Hum, D.W. 2002. Function of the transcriptional regulating protein of 132 kDa (TReP-132) on human P450scc gene expression. *Endocr. Res.* 28: 559-574.
- Gizard, F., Lavallee, B., DeWitte, F., Teissier, E., Staels, B. and Hum, D.W. 2002. The transcriptional regulating protein of 132 kDa (TReP-132) enhances P450scc gene transcription through interaction with steroidogenic factor-1 in human adrenal cells. *J. Biol. Chem.* 277: 39144-39155.
- Duguay, Y., Lapointe, A., Lavallee, B., Hum, D.W. and Rivest, S. 2003. Cloning of murine TReP-132, a novel transcription factor expressed in brain regions involved in behavioral and psychiatric disorders. *Mol. Psychiatry* 8: 39-49.
- Gizard, F., Teissier, E., Dufort, I., Luc, G., Luu-The, V., Staels, B. and Hum, D.W. 2004. The transcriptional regulating protein of 132 kDa (TReP-132) differentially influences steroidogenic pathways in human adrenal NCI-H295 cells. *J. Mol. Endocrinol.* 32: 557-569.
- Gizard, F., Robillard, R., Barbier, O., Quatannens, B., Faucompre, A., Revillion, F., Peyrat, J.P., Staels, B. and Hum, D.W. 2005. TReP-132 controls cell proliferation by regulating the expression of the cyclin-dependent kinase inhibitors p21^{WAF1/Cip1} and p27^{Kip1}. *Mol. Cell. Biol.* 25: 4335-4348.
- Gizard, F., Robillard, R., Gross, B., Barbier, O., Revillion, F., Peyrat, J.P., Torpier, G., Hum, D.W. and Staels, B. 2006. TReP-132 is a novel progesterone receptor coactivator required for the inhibition of breast cancer cell growth and enhancement of differentiation by progesterone. *Mol. Cell. Biol.* 26: 7632-7644.

CHROMOSOMAL LOCATION

Genetic locus: Trerf1 (mouse) mapping to 17 C.

PRODUCT

TReP-132 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TReP-132 shRNA Plasmid (m): sc-154633-SH and TReP-132 shRNA (m) Lentiviral Particles: sc-154633-V as alternate gene silencing products.

For independent verification of TReP-132 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154633A, sc-154633B and sc-154633C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TReP-132 siRNA (m) is recommended for the inhibition of TReP-132 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TReP-132 gene expression knockdown using RT-PCR Primer: TReP-132 (m)-PR: sc-154633-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.