



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



TRH-R2 siRNA (m): sc-154638

BACKGROUND

Thyrotrophin-releasing hormone (TRH) is a hypothalamic tripeptide that stimulates, via its receptor in the anterior pituitary gland, the release of thyrotrophin (TSH) and prolactin. The TRH receptors, TRH-R1 and TRH-R2, are G protein-coupled proteins containing seven transmembrane domains and other conserved regions. In rat, two isoforms exist, TRH-R (412) and TRH-R (387), that differ at their carboxy termini. TRH receptors are distributed throughout the central and peripheral nervous systems and are present in a variety of tissues. TRH-R2 displays 50% homology to TRH-R1 and is more restricted to the central nervous system than TRH-R1. Mutation in the TRH receptor gene is associated with isolated central hypothyroidism, a rare disorder characterized by insufficient TSH secretion resulting in low levels of thyroid hormones.

REFERENCES

- Eidne, K.A., et al. 1991. Cloning, sequencing and tissue distribution of a candidate G protein-coupled receptor from rat pituitary gland. *FEBS Lett.* 292: 243-248.
- de la Pena, P., et al. 1992. Two isoforms of the thyrotrophin-releasing hormone receptor generated by alternative splicing have indistinguishable functional properties. *J. Biol. Chem.* 267: 25703-25708.
- Zabavnik, J., et al. 1993. Distribution of thyrotrophin-releasing hormone receptor messenger RNA in rat pituitary and brain. *Neuroscience* 53: 877-887.
- Duthie, S.M., et al. 1993. Cloning and functional characterisation of the human TRH receptor. *Mol. Cell. Endocrinol.* 95: R11-R15.
- Cao, J., et al. 1998. Cloning and characterization of a cDNA encoding a novel subtype of rat thyrotrophin-releasing hormone receptor. *J. Biol. Chem.* 273: 32281-32287.
- Mitsuma, T., et al. 1999. Distribution of thyrotrophin releasing hormone receptor type 2 in rats: an immunohistochemical study. *Endocr. Regul.* 33: 135-139.
- Heuer, H., et al. 2000. Expression of thyrotrophin-releasing hormone receptor 2 (TRH-R2) in the central nervous system of rats. *J. Comp. Neurol.* 428: 319-336.
- O'Dowd, B.F., et al. 2000. TRH-R2 exhibits similar binding and acute signaling but distinct regulation and anatomic distribution compared with TRH-R1. *Mol. Endocrinol.* 14: 183-193.
- Collu, R. 2000. Genetic aspects of central hypothyroidism. *J. Endocrinol. Invest.* 23: 125-134.

CHROMOSOMAL LOCATION

Genetic locus: *Trhr2* (mouse) mapping to 8 E1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TRH-R2 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRH-R2 shRNA Plasmid (m): sc-154638-SH and TRH-R2 shRNA (m) Lentiviral Particles: sc-154638-V as alternate gene silencing products.

For independent verification of TRH-R2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154638A and sc-154638B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRH-R2 siRNA (m) is recommended for the inhibition of TRH-R2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRH-R2 gene expression knockdown using RT-PCR Primer: TRH-R2 (m)-PR: sc-154638-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.