



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TRP14 siRNA (m): sc-154689

BACKGROUND

Thioredoxins comprise a family of small proteins that, by catalyzing the oxidation of disulfide bonds, participate in redox reactions throughout the cell. Proteins that contain thioredoxin domains do not necessarily convey the oxidative properties of thioredoxins, but generally function as disulfide isomerases that enzymatically rearrange disulfide bonds found in various proteins. TRP14 (thioredoxin-related protein 14), also known as protein 42-9-9, TXNL5 or TXNDC17, is a 123 amino acid cytoplasmic protein that is ubiquitously expressed. Considered a disulfide reductase, TRP14 may participate in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions. TRP14 modulates TNF α signaling and NF κ B activation. Suggested to have peroxidase activity, TRP14 may contribute to the elimination of cellular hydrogen peroxide. TRP14 is reduced by TrXR1 and interacts with DYNLL1.

REFERENCES

1. Powis, G. and Montfort, W.R. 2001. Properties and biological activities of thioredoxins. *Annu. Rev. Biophys. Biomol. Struct.* 30: 421-455.
2. Jeong, W., Yoon, H.W., Lee, S.R. and Rhee, S.G. 2004. Identification and characterization of TRP14, a thioredoxin-related protein of 14 kDa. New insights into the specificity of thioredoxin function. *J. Biol. Chem.* 279: 3142-3150.
3. Jeong, W., Chang, T.S., Boja, E.S., Fales, H.M. and Rhee, S.G. 2004. Roles of TRP14, a thioredoxin-related protein in tumor necrosis factor α signaling pathways. *J. Biol. Chem.* 279: 3151-3159.
4. Woo, J.R., Kim, S.J., Jeong, W., Cho, Y.H., Lee, S.C., Chung, Y.J., Rhee, S.G. and Ryu, S.E. 2004. Structural basis of cellular redox regulation by human TRP14. *J. Biol. Chem.* 279: 48120-48125.
5. Nakamura, H. 2005. Thioredoxin and its related molecules: update 2005. *Antioxid. Redox Signal.* 7: 823-828.
6. Stefanková, P., Kollárová, M. and Barák, I. 2005. Thioredoxin-structural and functional complexity. *Gen. Physiol. Biophys.* 24: 3-11.

CHROMOSOMAL LOCATION

Genetic locus: Txndc17 (mouse) mapping to 11 B4.

PRODUCT

TRP14 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRP14 shRNA Plasmid (m): sc-154689-SH and TRP14 shRNA (m) Lentiviral Particles: sc-154689-V as alternate gene silencing products.

For independent verification of TRP14 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154689A, sc-154689B and sc-154689C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRP14 siRNA (m) is recommended for the inhibition of TRP14 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRP14 gene expression knockdown using RT-PCR Primer: TRP14 (m)-PR: sc-154689-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.