



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TSHZ2 siRNA (m): sc-154720

## BACKGROUND

The homeobox DNA-binding domain is a 60 amino acid motif that is conserved among many species and functions to bind DNA via a helix-turn-helix structure, thereby playing a role in transcriptional regulation and the control of gene expression. TSHZ2 (teashirt homolog 2), also known as ZNF218 (zinc finger protein 218) and OVC10-2 (ovarian cancer-related protein 10-2), is a 1,034 amino acid nuclear protein that contains one homeobox DNA-binding domain, as well as five C<sub>2</sub>H<sub>2</sub>-type zinc fingers. In *Drosophila*, teashirt genes function as region-specific homeotic genes that specify trunk identity during embryogenesis and these genes seem to conserve the same function in mice. The gene encoding TSHZ2 maps to human chromosome 20, which comprises approximately 2% of the human genome and contains nearly 63 million bases that encode over 600 genes.

## REFERENCES

1. Chauvet, S., et al. 2000. *dlarp*, a new candidate Hox target in *Drosophila* whose orthologue in mouse is expressed at sites of epithelium/mesenchymal interactions. *Dev. Dyn.* 218: 401-413.
2. Caubit, X., et al. 2000. Vertebrate orthologues of the *Drosophila* region-specific patterning gene teashirt. *Mech. Dev.* 91: 445-448.
3. Manfroid, I., et al. 2004. Three putative murine Teashirt orthologues specify trunk structures in *Drosophila* in the same way as the *Drosophila* teashirt gene. *Development* 131: 1065-1073.
4. Su, H.Y., et al. 2004. A novel gene homologous to teashirt is differentially expressed in neonatal mouse skin during development of hair follicles. *Mol. Biotechnol.* 28: 9-20.
5. Gocke, C.B., et al. 2005. Systematic identification and analysis of mammalian small ubiquitin-like modifier substrates. *J. Biol. Chem.* 280: 5004-5012.
6. Caubit, X., et al. 2005. Expression patterns of the three Teashirt-related genes define specific boundaries in the developing and postnatal mouse forebrain. *J. Comp. Neurol.* 486: 76-88.

## CHROMOSOMAL LOCATION

Genetic locus: *Tshz2* (mouse) mapping to 2 H3.

## PRODUCT

TSHZ2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TSHZ2 shRNA Plasmid (m): sc-154720-SH and TSHZ2 shRNA (m) Lentiviral Particles: sc-154720-V as alternate gene silencing products.

For independent verification of TSHZ2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154720A, sc-154720B and sc-154720C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

TSHZ2 siRNA (m) is recommended for the inhibition of TSHZ2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TSHZ2 gene expression knockdown using RT-PCR Primer: TSHZ2 (m)-PR: sc-154720-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.