



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TTC9C siRNA (m): sc-154784

BACKGROUND

The tetratricopeptide repeat (TPR) motif is a degenerate, 34 amino acid sequence found in many proteins and acts to mediate protein-protein interactions in various pathways. At the sequence level, there can be up to 16 tandem TPR repeats, each of which has a helix-turn-helix shape that stacks on other TPR repeats to achieve ligand binding specificity. TTC9C (tetratricopeptide repeat domain 9C) is a 171 amino acid protein belonging to the TTC9 family. Containing 3 TPR repeats, TTC9C is encoded by a gene located in a region of human chromosome 11q12.3, which houses over 1,400 genes and comprises nearly 4% of the human genome. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are associated with defects in genes that map to chromosome 11.

REFERENCES

- Blatch, G.L. and Lässle, M. 1999. The tetratricopeptide repeat: a structural motif mediating protein-protein interactions. *Bioessays* 21: 932-939.
- Andrade, M.A., Perez-Iratxeta, C. and Ponting, C.P. 2001. Protein repeats: structures, functions, and evolution. *J. Struct. Biol.* 134: 117-131.
- Smith, D.F. 2004. Tetratricopeptide repeat cochaperones in steroid receptor complexes. *Cell Stress Chaperones* 9: 109-121.
- Banerjee, A., Periyasamy, S., Wolf, I.M., Hinds, T.D., Yong, W., Shou, W. and Sanchez, E.R. 2008. Control of glucocorticoid and progesterone receptor subcellular localization by the ligand-binding domain is mediated by distinct interactions with tetratricopeptide repeat proteins. *Biochemistry* 47: 10471-10480.
- Wilson, J.B., Blom, E., Cunningham, R., Xiao, Y., Kupfer, G.M. and Jones, N.J. 2010. Several tetratricopeptide repeat (TPR) motifs of FANCG are required for assembly of the BRCA2/D1-D2-G-X3 complex, FANCD2 monoubiquitylation and phleomycin resistance. *Mutat. Res.* 689: 12-20.
- Schülke, J.P., Wochnik, G.M., Lang-Rollin, I., Gassen, N.C., Knapp, R.T., Berning, B., Yassouridis, A. and Rein, T. 2010. Differential impact of tetratricopeptide repeat proteins on the steroid hormone receptors. *PLoS ONE* 5: e11717.
- Krachler, A.M., Sharma, A. and Kleanthous, C. 2010. Self-association of TPR domains: lessons learned from a designed, consensus-based TPR oligomer. *Proteins* 78: 2131-2143.

CHROMOSOMAL LOCATION

Genetic locus: *Ttc9c* (mouse) mapping to 19 A.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TTC9C siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TTC9C shRNA Plasmid (m): sc-154784-SH and TTC9C shRNA (m) Lentiviral Particles: sc-154784-V as alternate gene silencing products.

For independent verification of TTC9C (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154784A, sc-154784B and sc-154784C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TTC9C siRNA (m) is recommended for the inhibition of TTC9C expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TTC9C gene expression knockdown using RT-PCR Primer: TTC9C (m)-PR: sc-154784-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.