



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TLL5 siRNA (m): sc-154793

## BACKGROUND

A large protein group known as the Tubulin tyrosine ligase-like (TLL) family is implied to catalyze ligations of amino acids to Tubulins and other substrates. Each member contains a characteristic TTL domain. TLL5 (Tubulin tyrosine ligase-like family, member 5), also known as STAMP (SRC1 and TIF2-associated modulatory protein), is a 1,281 nuclear and cytoplasmic protein that belongs to the Tubulin-tyrosine ligase family. Containing one TTL domain, TLL5 is highly expressed in heart and skeletal muscle and lowly expressed in other tissues. TLL5 plays a role as a Tubulin polyglutamylase and participates in forming polyglutamate side chains on Tubulin. TLL5 interacts with and magnifies the effect of transcriptional coactivator GRIP-1, thereby regulating glucocorticoid receptor-mediated repression and induction as well as in androgen receptor-mediated induction. TLL5 exists as three alternatively spliced isoforms and is encoded by a gene located on human chromosome 14q24.3.

## REFERENCES

1. Trichet, V., Ruault, M., Roizès, G. and De Sario, A. 2000. Characterization of the human Tubulin tyrosine ligase-like 1 gene (TLL1) mapping to 22q13.1. *Gene* 257: 109-117.
2. Janke, C., Rogowski, K., Wloga, D., Regnard, C., Kajava, A.V., Strub, J.M., Temurak, N., van Dijk, J., Boucher, D., van Dorsselaer, A., Suryavanshi, S., Gaertig, J. and Edde, B. 2005. Tubulin polyglutamylase enzymes are members of the TTL domain protein family. *Science* 308: 1758-1762.
3. He, Y. and Simons, S.S., Jr. 2007. STAMP, a novel predicted factor assisting TIF2 actions in glucocorticoid receptor-mediated induction and repression. *Mol. Cell. Biol.* 27: 1467-1485.
4. van Dijk, J., Miro, J., Strub, J.M., Lacroix, B., van Dorsselaer, A., Edde, B. and Janke, C. 2008. Polyglutamylation is a post-translational modification with a broad range of substrates. *J. Biol. Chem.* 283: 3915-3922.
5. Sun, Y., Tao, Y.G., Kagan, B.L., He, Y. and Simons, S.S., Jr. 2008. Modulation of transcription parameters in glucocorticoid receptor-mediated repression. *Mol. Cell. Endocrinol.* 295: 59-69.
6. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612268. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. He, Y., Blackford, J.A., Jr., Kohn, E.C. and Simons, S.S., Jr. 2010. STAMP alters the growth of transformed and ovarian cancer cells. *BMC Cancer* 10: 128.

## CHROMOSOMAL LOCATION

Genetic locus: Tll5 (mouse) mapping to 12 D2.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

TLL5 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TLL5 shRNA Plasmid (m): sc-154793-SH and TLL5 shRNA (m) Lentiviral Particles: sc-154793-V as alternate gene silencing products.

For independent verification of TLL5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154793A, sc-154793B and sc-154793C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

TLL5 siRNA (m) is recommended for the inhibition of TLL5 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contains a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TLL5 gene expression knockdown using RT-PCR Primer: TLL5 (m)-PR: sc-154793-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.