



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TXA2R siRNA (m): sc-154816

BACKGROUND

Thromboxane A2 (TXA2), the major cyclooxygenase (COX) product of arachidonic acid, stimulates platelet aggregation and is a potent vasoconstrictor. TXA2R has been implicated in several processes in normal kidney physiology as well as in myocardial infarction, atherosclerosis and bronchial asthma. TXA2 mediates its effects through the TXA2 receptor (TXA2R), a G protein-coupled receptor that activates phospholipase C (PLC) to mobilize intracellular calcium stores. Alternatively, TXA2R activates the MAP kinase pathway in response to dibutyl cyclic AMP (dbcAMP). The human TXA2R gene maps to chromosome 19p13.3 and is alternatively spliced to produce proteins which differ in the carboxy-termini. TXA2R is expressed in platelets, endothelium, placenta, vascular smooth muscles and the renal cortex. Mutations in the gene encoding TXA2R lead to several bleeding disorders due to either impaired coupling between the receptor and PLC or impaired binding to TXA2. Also, the TXA2R protein may be involved in mediating renal damage in disease states, controlling the initiation and/or progression of labor in women, and providing therapeutic value for treatment of acute pancreatitis.

REFERENCES

- Swanson, M.L., et al. 1992. The expression of thromboxane A2 synthase and thromboxane A2 receptor gene in human uterus. *Biol. Reprod.* 47: 105-117.
- Reilly, M., et al. 1993. Cellular activation by thromboxane A2 and other eicosanoids. *Eur. Heart J.* 14: 88-93.
- Nusing, R.M., et al. 1993. Characterization and chromosomal mapping of the human thromboxane A2 receptor gene. *J. Biol. Chem.* 268: 25253-25259.
- Raychowdhury, M.K., et al. 1994. Alternative splicing produces a divergent cytoplasmic tail in the human endothelial thromboxane A2 receptor. *J. Biol. Chem.* 269: 19256-19261.
- Hirata, T., et al. 1996. Two thromboxane A2 receptor isoforms in human platelets. Opposite coupling to adenylyl cyclase with different sensitivity to Arg60 to Leu mutation. *J. Clin. Invest.* 97: 949-956.

CHROMOSOMAL LOCATION

Genetic locus: Tbx2r (mouse) mapping to 10 C1.

PRODUCT

TXA2R siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TXA2R shRNA Plasmid (m): sc-154816-SH and TXA2R shRNA (m) Lentiviral Particles: sc-154816-V as alternate gene silencing products.

For independent verification of TXA2R (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154816A and sc-154816B.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TXA2R siRNA (m) is recommended for the inhibition of TXA2R expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TXA2R gene expression knockdown using RT-PCR Primer: TXA2R (m)-PR: sc-154816-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.