



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TYW4 siRNA (m): sc-154830

BACKGROUND

Wybutosine, a derivative of wyosine, is a tricyclic hypermodified guanosine found in eukaryotic and archaeal tRNAs. TYW4 (tRNA-γW synthesizing protein 4), also known as PPM2 (p21WAF1/CIP1 promoter-interacting protein) or LCMT2 (leucine carboxyl methyltransferase 2), is a 686 amino acid protein belonging to the methyltransferase superfamily. A component of the wybutosine (γW) biosynthesis pathway, TYW4 may act as a S-adenosyl-L-methionine-dependent methyltransferase that catalyzes the final step of γW biosynthesis, methylation and methoxycarbonylation. TYW4 is encoded by a gene located on human chromosome 15, which houses over 700 genes and comprises nearly 3% of the human genome. Angelman syndrome, Prader-Willi syndrome, Tay-Sachs disease and Marfan syndrome are all associated with defects in chromosome 15-localized genes.

REFERENCES

1. Noma, A., Kirino, Y., Ikeuchi, Y. and Suzuki, T. 2006. Biosynthesis of wybutosine, a hyper-modified nucleoside in eukaryotic phenylalanine tRNA. *EMBO J.* 25: 2142-2154.
2. Noma, A. and Suzuki, T. 2006. Ribonucleome analysis identified enzyme genes responsible for wybutosine synthesis. *Nucleic Acids Symp. Ser.* 50: 65-66.
3. Zody, M.C., Garber, M., Sharpe, T., Young, S.K., Rowen, L., O'Neill, K., Whittaker, C.A., Kamal, M., Chang, J.L., Cuomo, C.A., Dewar, K., Fitzgerald, M.G., Kodira, C.D., Madan, A., Qin, S., Yang, X., Abbasi, N., et al. 2006. Analysis of the DNA sequence and duplication history of human chromosome 15. *Nature* 440: 671-675.
4. Makoff, A.J. and Flomen, R.H. 2007. Detailed analysis of 15q11-q14 sequence corrects errors and gaps in the public access sequence to fully reveal large segmental duplications at breakpoints for Prader-Willi, Angelman, and inv dup(15) syndromes. *Genome Biol.* 8: R114.
5. Suzuki, Y., Noma, A., Suzuki, T., Senda, M., Senda, T., Ishitani, R. and Nureki, O. 2007. Crystal structure of the radical SAM enzyme catalyzing tricyclic modified base formation in tRNA. *J. Mol. Biol.* 372: 1204-1214.
6. Goto-Ito, S., Ishii, R., Ito, T., Shibata, R., Fusatomi, E., Sekine, S.I., Bessho, Y. and Yokoyama, S. 2007. Structure of an archaeal TYW1, the enzyme catalyzing the second step of wye-base biosynthesis. *Acta Crystallogr. D Biol. Crystallogr.* 63: 1059-1068.
7. Suzuki, Y., Noma, A., Suzuki, T., Ishitani, R. and Nureki, O. 2009. Structural basis of tRNA modification with CO₂ fixation and methylation by wybutosine synthesizing enzyme TYW4. *Nucleic Acids Res.* 37: 2910-2925.

CHROMOSOMAL LOCATION

Genetic locus: *Lcmt2* (mouse) mapping to 2 E5.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TYW4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TYW4 shRNA Plasmid (m): sc-154830-SH and TYW4 shRNA (m) Lentiviral Particles: sc-154830-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TYW4 siRNA (m) is recommended for the inhibition of TYW4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TYW4 gene expression knockdown using RT-PCR Primer: TYW4 (m)-PR: sc-154830-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.